

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 0 896 002 A1**

(12)

**EUROPEAN PATENT APPLICATION**

published in accordance with Art. 158(3) EPC

(43) Date of publication:

10.02.1999 Bulletin 1999/06

(51) Int. Cl.<sup>6</sup>: **C07K 14/705**, **C12N 15/12**,  
**G01N 33/50**

(21) Application number: **98901043.4**

(86) International application number:  
**PCT/JP98/00370**

(22) Date of filing: **29.01.1998**

(87) International publication number:  
**WO 98/32771 (30.07.1998 Gazette 1998/30)**

(84) Designated Contracting States:  
**DE FR GB IT**

(30) Priority: **29.01.1997 JP 15118/97**  
**29.08.1997 JP 234544/97**

(71) Applicant:  
**TORAY INDUSTRIES, INC.**  
**Tokyo 103 (JP)**

(72) Inventors:  
• **KAINOH, Mie**  
**Kanagawa 251 (JP)**  
• **TANAKA, Toshiaki**  
**kanagawa 249 (JP)**

(74) Representative: **Kador & Partner**  
**Corneliusstrasse 15**  
**80469 München (DE)**

(54) **CHIMERIC PROTEINS, HETERODIMER COMPLEXES THEREOF AND SUBSTITUTE FOR PLATELET**

(57) Integrin-immunoglobulin chimeric protein heterodimer complexes in which the  $\alpha$ -chain and  $\beta$ -chain of integrins have been associated in a stable state. These complexes are not only usable as drugs as such but also applicable to the assaying of binding of integrins to ligands and the detection of substances binding to integrins or those inhibiting binding of integrins to ligands. These complexes are also usable as diagnostic agents. It has been found that integrins isolated in a stably associated structure would bind to extracellular matrixes under physiological conditions in the presence of plasma components, which indicates that integrins and, in its turn, extracellular matrix receptors might be usable as substitutes for platelets.

**EP 0 896 002 A1**

**BEST AVAILABLE COPY**

## Description

## Technical Field:

- 5 [0001] The present invention relates to chimeric proteins consisting of an integrin and an immunoglobulin, their heterodimer complexes, a production process thereof, their applications as drugs and reagents, etc. Furthermore, the present invention relates to medicinal application of isolated extracellular matrix receptors such as integrin-immunoglobulin chimeric protein heterodimer complexes, as platelet substitutes.

## 10 Background Arts:

[0002] Various cells have receptors which mediate the adhesion between a cell and a cell and receptors which mediate the adhesion between a cell and an extracellular matrix, and these receptors play important roles in immune reaction, inflammatory reaction, development, morphogenesis, wound healing, hemostasis, cancerous metastasis, etc. By separating and identifying the receptors which participate in these phenomena, the existence of so-called cell adhesion molecules has been clarified. Many of the molecules identified one after another are classified in reference to their structural features into integrin superfamily, immunoglobulin superfamily, selectin family, cadherin family, etc. (Corliss, T. M. and Harlan, J. M., Blood, 84, 2068-2101 (1994)). Of these families, the immunoglobulin superfamily, selectin family and cadherin family mediate mainly the adhesion between a cell and a cell, while the integrin superfamily is the so-called extracellular matrix receptors which mediate the adhesion to extracellular matrices such as fibronectin and collagens. In addition, extracellular matrix receptors which do not belong to any of these adhesion molecule families include CD26 (DDP1V), CD44, GPIV, GPVI, GPIb-vWF, etc. CD26 is a receptor for collagens, and CD44 is a receptor for hyaluronic acid, fibronectin and collagens ("Adhesion Molecules" p. 32-42, Masayuki Miyasaka (1991), Medical View (in Japanese)). Furthermore, it is reported that among the membrane glycoproteins (GPs) existing on platelets, GPIV, GPVI, GPIb-vWF, etc. are also collagen receptors ("Platelet Receptors", p. 119-132, Minoru Ohkuma et al., (1992), Kin-podo (in Japanese)).

[0003] A receptor belonging to the integrin superfamily has a heterodimer complex structure in which two subunits,  $\alpha$ -chain and  $\beta$ -chain as mutually different membrane proteins are associated with each other non-covalently (Hynes, R. O., Cell, 48, 549-554 (1987)). In the past, the integrin superfamily was classified into three subfamilies;  $\beta$ 1 integrin,  $\beta$ 2 integrin and  $\beta$ 3 integrin. Later, new  $\beta$  chains and  $\alpha$  chains were discovered one after another, and presently eight  $\beta$  chains ( $\beta$ 1,  $\beta$ 2,  $\beta$ 3,  $\beta$ 4,  $\beta$ 5,  $\beta$ 6,  $\beta$ 7 and  $\beta$ 8 and fifteen  $\alpha$  chains ( $\alpha$ 1,  $\alpha$ 2,  $\alpha$ 3,  $\alpha$ 4,  $\alpha$ 5,  $\alpha$ 6,  $\alpha$ 7,  $\alpha$ 8,  $\alpha$ 9,  $\alpha$ v,  $\alpha$ L,  $\alpha$ M,  $\alpha$ X,  $\alpha$ 1b and  $\alpha$ E) have been identified (Elner, S. G. and Elner, V. M., Inv. Ophtal. Vis. Sci., 37, 696-701 (1996)). It is known that each  $\beta$  chain is associated with one to eight  $\alpha$  chains, and as a result, 21 pairs of an  $\alpha$  chain and a  $\beta$  chain, i.e., integrin molecules have been identified (Elner, S. G. and Elner, V. M., Inv. Ophtal. Vis. Sci., 37, 696-701 (1996)). They include  $\alpha$ 4 $\beta$ 1 (VLA-4,  $\beta$ 1 integrin),  $\alpha$ L $\beta$ 2 (LFA-1,  $\beta$ 2 integrin),  $\alpha$ M $\beta$ 2 (Mac-1,  $\beta$ 2 integrin),  $\alpha$ 1b $\beta$ 3 (GPIIb/IIIa,  $\beta$ 3 integrin), etc. now targeted for drug development (Drug and Market Development, 6, 201-205 (1995)). Many other integrins are also expected to have relations with diseases.

[0004] The heterodimer complex structure of an integrin plays an important role in binding to a ligand (Hynes, R. O., Cell, 48, 549-554 (1987)). For example, it is estimated that the ligand binding region on an integrin consists of both an  $\alpha$  chain and a  $\beta$  chain (Hynes, R. O., Cell, 69, 11-25 (1992)). The fact that integrins having the same  $\alpha$  chain but associated with a different  $\beta$  chain, or integrins having the same  $\beta$  chain but associated with a different  $\alpha$  chain are respectively different in substrate specificity (Elner, S. G. and Elner, V. M., Inv. Ophtal. Vis. Sci. 37, 696-701 (1996)) supports this assumption. On the other hand, it was reported that the  $\alpha$  chains of some integrins have an sequence called an I domain consisting of about 180 amino acids inserted in the molecule, and data suggest that the I domain only could be bound to a ligand were reported (Ueda, T. et al., Proc. Natl. Acad. Sci. USA, 91, 10680-10684 (1994)). However, it was also reported that the I domain of an  $\alpha$  domain and the integrin as its original heterodimer complex are different in the style of binding to a ligand (Kamata, T. and Takada, Y., J. Biol. Chem., 269, 26006-26010 (1994)). It is also not clarified yet whether such parameters as specificity and affinity to a ligand are identical. It is not reported that in the case of an integrin not containing the I domain, for example, in the case of  $\alpha$ 4 $\beta$ 1 a partial structure only is bound to a ligand.

50 [0005] If any integrin isolated and prepared retains its heterodimer complex structure, hence the ligand binding capability, it can be used for studying the style of binding to a ligand in a state close to nature. Furthermore, it can be used as it is as a drug and can also be used as a reagent for measuring the amount of a ligand in tissue or serum or as a material for searching for adhesion inhibiting compounds very usefully. However, isolating and preparing an integrin with its function retained is said to be very difficult. One reason is that since the association between an  $\alpha$  chain and a  $\beta$  chain of an integrin is maintained non-covalently as described before, they are easily dissociated during isolation and preparation. Since an integrin is a membrane protein, the necessity of using a surfactant, etc. for solubilization is considered to be a large cause in the dissociation of the complex. In other words, the non-covalent preservation of functional structure inhibits the preparation of such an integrin.

[0006] In spite of the difficulty as described above, some cases were reported, in which an integrin heterodimer complex was isolated and prepared with its function retained. For cases of  $\alpha 2\beta 1$ ,  $\alpha 5\beta 1$  and  $\alpha v\beta 3$ , it was reported that the binding to a ligand can be determined by letting a liposome incorporate an integrin purified by using affinity column chromatography (Santoro, S. A. et al., *Biochem. Biophys. Res. Comm.*, 153, 217-223 (1988), Pytela, R. et al. *Cell*, 40, 191-198 (1985), Pytela, R. et al., *Method Enzymol.*, 144, 475-489 (1987)). For other cases, it was that if purified  $\alpha 5\beta 1$  or  $\alpha v\beta 3$  is coated on a plate, a peptide which inhibits the cell adhesion through the integrin can be selected (Koivunen, E. et al., *J. Biol. Chem.*, 268, 20205-20210 (1993), Healy, J. M. et al., *Biochemistry*, 34, 3948-3955 (1995)). For further other cases, it was reported that if purified  $\alpha v\beta 3$  or  $\alpha 4\beta 1$  is coated on a plate, the binding to a ligand can be determined (Charo, I. F. et al., *J. Cell Biol.*, 111, 2795-2800 (1990), Makarew, R. et al., *J. Biol. Chem.*, 269, 4005-4011 (1994), Paul Mould, A. et al., *J. Biol. Chem.*, 269, 27224-27230 (1994)). For a still further other case, it was reported that if an extracellular portion of  $\alpha 1\beta 3$  heterodimer complex prepared by gene manipulation is coated on a plate through a complex specific antibody, the binding to a ligand can be determined (Gulino, D. et al., *Eur. J. Biochem.*, 227, 108-115 (1995)). These cases suggest that to exert the function of a purified integrin, its heterodimer complex must be bound to or included in any carrier. The reason why a carrier is considered to be necessary is that since a heterodimer complex is associated non-covalently in a solution, it tends to be dissociated and as a result, cannot retain its functional structure. In the finally stated case, only a molecule with a heterodimer complex structure is selected using a complex specific antibody, in a design to determine the binding in a state where both the chains are not dissociated from each other.

[0007] As a case requiring no carrier, it was reported that purified  $\alpha 1\beta 1$ , or  $\alpha 2\beta 1$  allows the determination of the binding to a ligand dependent on high concentration of metal ions even without using any carrier (Pfaff, M. et al., *Eur. J. Biochem.*, 225, 975-984 (1994)). In this case, the surfactant added in the process of purification plays a role similar to that of a liposome, acting as a carrier. For a further other case, it was reported that an extracellular of  $\alpha M\beta 2$  heterodimer complex prepared by using gene manipulation is bound to a ligand (Berman, P. W. et al., *J. Cell Biochem.*, 52, 183-195 (1993)). These cases do not suggest the necessity of any carrier as described before, but the disadvantage that the association of molecules in a heterodimer complex is retained non-covalently is not improved.

[0008] As a still further other case, a chimeric protein consisting of  $\alpha d$  and an immunoglobulin is disclosed (Japanese Patent Laid-Open (Kokai) No. 8-507933), but only the result of immune precipitation is reported, without examining the binding to a ligand. Furthermore, since a  $\beta$  chain is not expressed in the chimeric protein as an immunoglobulin, the binding between an  $\alpha$  chain and a  $\beta$  chain remains non-covalent.

[0009] The above facts suggest that any integrin with an  $\alpha$  chain and a  $\beta$  chain structurally stably associated and with its function retained has never been successfully prepared. That a complex structure is unstable restricts the use of its molecule.

[0010] Of the molecules belonging to the integrin superfamily, integrin  $\alpha 2\beta 1$  is an extracellular matrix receptor found to be expressed in T cells, platelets, etc. activated for long time. However, it was reported that the  $\alpha 2\beta 1$  on the cell surfaces of platelets and fibroblasts is bound to collagens only and that the  $\alpha 2\beta 1$  on the surfaces of vascular endothelial cells is bound to both collagens and laminins (Elices, M. J. et al., *Proc. Natl. Acad. Sci. USA*, 86, 9906-9910 (1989)), and it is speculated that the function of  $\alpha 2\beta 1$  becomes different, depending on cells.

[0011] In relation to the conditions of diseases, there are reports to suggest that integrin  $\alpha 2\beta 1$  plays an important role for wound healing and cancerous metastasis (Shiro, J. A. et al., *Cell*, 67, 403-410 (1991), Chen, F. et al., *J. Exp. Med.*, 173, 1111-1119 (1991), Chan, B. M. C. et al., *Science*, 251, 1600-1602 (1991)). Furthermore, it was reported that from the analysis of platelet function of patients with bleeding tendency, the adherence of platelets and collagens through integrin  $\alpha 2\beta 1$  has close relation with the first step of hemostasis and thrombosis process (Nieuwenhuis, H. K. et al., *Nature*, 318, 470-472 (1985)). Though the relations of integrin  $\alpha 2\beta 1$  with conditions of diseases are suggested like this, any medical application of using the integrin  $\alpha 2\beta 1$  protein and other isolated extracellular matrix receptor proteins under physiological ion condition or in the presence of plasma components has not been examined.

[0012] On the other hand, the necessity for artificial substitutes of platelets used as blood preparations in the clinical field is growing, and various attempts have been reported (*Progress of Medicine* 179, 406-407 (1996), *Clinical Blood* 37, 1353-1361 (1997) (respectively in Japanese)). However, they are not yet practically available.

#### Disclosure of the Invention:

[0013] The present invention relates to chimeric proteins in which the  $\alpha$  chain and  $\beta$  chain of an integrin are combined with the heavy chain or light chain of an immunoglobulin, their heterodimer complexes, a production process thereof, a method for testing the binding of an integrin-immunoglobulin chimeric protein heterodimer complex to a ligand and a cell, substances bound to an integrin obtained by using the method, a method for searching for a substance inhibiting the binding between an integrin and a ligand using the integrin-immunoglobulin chimeric protein heterodimer complex, substances for inhibiting the binding, and the application of integrin-immunoglobulin chimeric protein heterodimer complexes as drugs and reagents. Furthermore, the present invention relates to platelet substitutes containing an integrin-immunoglobulin chimeric protein heterodimer complex or any other isolated extracellular matrix receptor as an active

ingredient.

# Brief Description of the Drawings:

5 [0014]

Fig. 1 shows that  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex is bound to VCAM-1 expressing cell, and that the binding is inhibited by an anti-integrin antibody or EDTA, a cationic cheating agent.

10 Fig. 2 shows that  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex is bound to CS-1 peptide, and that the binding is inhibited by an anti-integrin antibody or EDTA, a cationic cheating agent.

Fig. 3 shows that the binding between  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1$  IgG heavy chain chimeric protein heterodimer complex and CS-1 peptide is inhibited by GPEILDVPST, and is not inhibited by any other peptide.

15 Fig. 4 shows that  $\alpha 2 \cdot$  IgG heavy chain- $\beta 1$  IgG heavy chain chimeric protein heterodimer complex is bound to a collagen, and that the binding is inhibited by an anti-integrin antibody and EDTA, a cationic cheating agent.

Fig. 5 shows that  $\alpha 2 \cdot$  IgG heavy chain- $\beta 1$  IgG heavy chain chimeric protein heterodimer complex liposome is bound to a collagen in the presence of plasma.

Fig. 6 shows that the binding of  $\alpha 2 \cdot$  IgG heavy chain- $\beta 1$  IgG heavy chain chimeric protein heterodimer complex liposome to a collagen is inhibited by an anti-integrin antibody or EDTA, a cationic chelating agent.

20

## The Best Embodiments of the Invention:

[0015] The extracellular matrix receptors in the present invention refer generally to the receptors which mediate the adhesion between a cell and an extracellular matrix. The receptors include the integrin superfamily having a heterodimer complex structure in which an  $\alpha$  chain and a  $\beta$  chain are non-covalently associated with each other as two membrane proteins (Corliss, T. M. and Harlan, J. M. Blood, 84, 2068-2101 (1994)), and other receptors such as CD26 (DDP1V), CD44, GPIV, GPVI, GPb-vWF, etc. The integrins in the present invention refer to molecules belonging to the integrin superfamily, and also include the isomers of the molecules belonging to the family. The  $\alpha$  chains of the present invention include 15  $\alpha$  chains, i.e.,  $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 3$ ,  $\alpha 4$ ,  $\alpha 5$ ,  $\alpha 6$ ,  $\alpha 7$ ,  $\alpha 8$ ,  $\alpha 9$ ,  $\alpha v$ ,  $\alpha L$ ,  $\alpha M$ ,  $\alpha X$ ,  $\alpha 11b$  and  $\alpha E$ , and among them,  $\alpha 4$  and  $\alpha 2$  are preferable, though preferable  $\alpha$  chains are not limited to them. The  $\beta$  chains of the present invention include eight  $\beta$  chains, i.e.,  $\beta 1$ ,  $\beta 2$ ,  $\beta 3$ ,  $\beta 4$ ,  $\beta 5$ ,  $\beta 6$ ,  $\beta 7$  and  $\beta 8$  and among them,  $\beta 1$  is preferable, though preferable  $\beta$  chains are not limited to it. The integrin molecules as pairs respectively consisting of an  $\alpha$  chain and a  $\beta$  chain include the twenty one integrins stated in Elnar, S. G. and Elnar, V. M., Inv. Ophthal. Vis. Sci., 37, 696-701 (1996), though not limited to them.

35 [0016] A chimeric protein consisting of the  $\alpha$  chain of an integrin and the heavy chain or light chain of an immunoglobulin refers to a molecule in which the extracellular region of the  $\alpha$  chain of an integrin is bound to the constant region of the heavy chain or light chain contained an immunoglobulin. In this case, a chimeric protein in which N terminus side of the protein is integrin molecule and then connected to an immunoglobulin molecule side by side is preferable. A chimeric protein consisting of the  $\beta$  chain of an integrin and the heavy chain or light chain of an immunoglobulin refers to a molecule in which the extracellular region of the  $\beta$  chain of an integrin is bound to the constant region of the heavy chain or light chain contained in an immunoglobulin. Also in this case, a chimeric protein in which N terminus side of the protein is an integrin molecule and then connected to an immunoglobulin molecule side by side is preferable. In either case of  $\alpha$  chain or  $\beta$  chain, a chimeric protein bound to the heavy chain of an immunoglobulin is preferable.

40 [0017] The isotype of the immunoglobulin to be bound to the  $\alpha$  chain or  $\beta$  chain is not especially limited. Any of IgG, IgM, IgA and IgE can be used, but it is preferable to use IgG. The subclasses of IgG include IgG<sub>1</sub>, IgG<sub>2</sub>, IgG<sub>3</sub> and IgG<sub>4</sub>, but it is preferable to use IgG<sub>1</sub>. Furthermore, it is possible to use a molecule with a dimer structure having a disulfide bond between molecules instead of the immunoglobulin.

[0018] In the present invention, a molecule in which a chimeric protein consisting of the  $\alpha$  chain of an integrin and the heavy chain or light chain of an immunoglobulin and a chimeric protein consisting of the  $\beta$  chain of the integrin and the heavy chain or light chain of the immunoglobulin are associated with each other is called an integrin-immunoglobulin chimeric protein heterodimer complex. In this case, a combination consisting of  $\alpha$  chain  $\cdot$  immunoglobulin heavy chain (which means a chimeric protein consisting of an  $\alpha$  chain and the heavy chain of an immunoglobulin; hereinafter this applies) and  $\beta$  chain  $\cdot$  immunoglobulin heavy chain, a combination consisting of  $\alpha$  chain  $\cdot$  immunoglobulin heavy chain and  $\beta$  chain  $\cdot$  immunoglobulin light chain, and a combination consisting of  $\alpha$  chain  $\cdot$  immunoglobulin light chain and  $\beta$  chain  $\cdot$  immunoglobulin heavy chain are preferable. A combination consisting of  $\alpha$  chain  $\cdot$  immunoglobulin heavy chain and  $\beta$  chain  $\cdot$  immunoglobulin heavy chain is more preferable.

55 [0019] In the integrin-immunoglobulin chimeric protein heterodimer complex of the present invention, the  $\alpha$  chain can be  $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 3$ ,  $\alpha 4$ ,  $\alpha 5$ ,  $\alpha 6$ ,  $\alpha 7$ ,  $\alpha 8$ ,  $\alpha 9$ ,  $\alpha v$ ,  $\alpha L$ ,  $\alpha M$ ,  $\alpha X$ ,  $\alpha 11b$  or  $\alpha E$ , and the  $\beta$  chain can be  $\beta 1$ ,  $\beta 2$ ,  $\beta 3$ ,  $\beta 4$ ,  $\beta 5$ ,  $\beta 6$ ,  $\beta 7$  or

$\beta 8$ , It is preferable that the  $\alpha$  chain is  $\alpha 4$  or  $\alpha 2$  and that the  $\beta$  chain is  $\beta 1$ , though preferable chains are not limited to them.

[0020] The process for preparing an integrin-immunoglobulin chimeric protein heterodimer complex is described below, but the process is not limited thereto.

5 [0021] A DNA coding for the  $\alpha$  chain and  $\beta$  chain of an integrin can be obtained using the information of known cDNA sequences by such a method as gene amplification based on the PCR method, cDNA cloning or DNA synthesis. For example, the DNA sequences of  $\alpha 4$  and  $\beta 1$  are already reported in literature (Takada, Y. et al., EMBO J., 8, 1361-1368 (1989), Scott Graves, W. et al., J. Cell Biol., 105, 1183-1190 (1987)). A DNA coding for the  $\alpha$  chain and  $\beta$  chain of an integrin can also be obtained by the expression cloning using an antibody, etc. For binding to a DNA coding for the constant region of an immunoglobulin, it is desirable to take out a DNA coding for the extracellular portions only of the  $\alpha$  chain and  $\beta$  chain of an integrin. For this purpose, it is preferable to use the PCR method and DNA synthesis. The extracellular portion of either an  $\alpha$  chain or  $\beta$  chain refers to the polypeptide sequence on the N terminus side from the portion speculated to be the transmembrane portion. Its partial sequence can also be used as far as the ligand binding capability is retained, but it is preferable to use most of the portion considered to be an extracellular region. For taking out a DNA, it is necessary to adjust for adaptation of frames after linking to a DNA coding for an immunoglobulin. For example, this can be achieved by modifying the primer when a DNA fragment is taken out by the PCR method. In this case, it is desirable to design for ensuring that amino acid modification is not caused by the base substitution of the primer. However, amino acid substitution is allowed as far as the function of the chimeric protein is not changed. For obtaining a DNA by chemical synthesis, the purpose can be achieved by designing a sequence to ensure the linking to a DNA coding for an immunoglobulin. In the case of cDNA, a DNA capable of being bound to a DNA coding for an immunoglobulin can be prepared by using DNA fragmentation and a synthetic DNA.

15 [0022] Then, a DNA coding for an immunoglobulin is prepared. In the present invention, it is desirable to use DNAs coding for the heavy chain and light chain of a human immunoglobulin, but DNAs coding for an immunoglobulin of another animal species can also be used. The preparation of a DNA coding for human IgG is already reported (Ellison, J. W. et al., Nucleic Acids Res., 10, 4071-4079 (1982)), but the preparation is not limited to this method. Any method similar to the above mentioned method for preparing DNAs coding for the  $\alpha$  chain and  $\beta$  chain of an integrin can also be used. In the present invention, for the heavy chain of a human immunoglobulin, it is preferable to use a genomic DNA, but a cDNA can also be used. As the DNA for the heavy chain of a human immunoglobulin, it is preferable to use a portion coding for the hinge region, CH2 region or CH3 region, but a DNA coding for the entire constant region of CH1 or CH3 can also be used. For the light chain of an immunoglobulin, a DNA coding for the CL region is used. Finally, a DNA coding for the extracellular portion of an  $\alpha$  chain or  $\beta$  chain and a DNA coding for the constant region of human immunoglobulin heavy chain are linked with in frame. The obtained DNA codes for a polypeptide starting from the methionine of translation initiation and having the signal sequence of the  $\alpha$  chain or  $\beta$  chain of an integrin, its extracellular region and the constant region of human immunoglobulin heavy chain linked in this order.

25 [0023] The DNA coding for a chimeric protein consisting of the  $\alpha$  chain of an integrin and the heavy chain or light chain of an immunoglobulin, or the DNA coding for a chimeric protein consisting of the  $\beta$  chain of an integrin and the heavy chain or light chain of an immunoglobulin respectively obtained in the above is functionally linked in a proper expression control sequence, to obtain a recombinant vector. The general methods concerning gene recombination such as the method for preparing the recombinant vector, the method for transfecting it into a cell are described in a published book ("Molecular Cloning", Sambrook et al., (1989) Cold Spring Harbor Lab. Press, New York), but the methods are not limited to those stated there. In the present invention, it is desirable to use an expression control sequence suitable for protein expression in animal cells. For example, for manifestation of insect cells, polyhedrin promoter, p10 promoter, etc. are generally used as expression control sequences, and for expression of other animals' cells, SR $\alpha$  promoter, cytomegalovirus derived promoter, simian virus 40 derived promoter, polyhedrin promoter, p10 promoter, etc. are used. However, the expression control sequences are not limited to them. In the present invention, it is preferable to use SR $\alpha$  promoter.

35 [0024] If the obtained recombinant vector is transfecting into a cell, a cell capable of producing an integrin-immunoglobulin chimeric protein heterodimer complex can be obtained. In this case it is preferable to use an animal derived cell as a host. For example, COS cell (simian renal cell), CHO cell (Chinese Hamster ovarian cell), Sf9 (insect cell), etc. are generally used as hosts. Furthermore, myeloma cells such as P3U1 and Y3 can also be used. Other established cell lines and cloned cells can also be used, but the cells used as hosts are not limited to them. In the present invention, it is preferable to use a CHO cell.

40 [0025] It is known that the methods for transfecting a recombinant vector into a cell include the lipofectin method, calcium phosphate method, electroporation method, etc., and any of the methods can be used. The method is not limited to them. It is preferable that when a cell is transfecting by using a recombinant vector, a recombinant vector for expression of a chimeric protein consisting of the  $\alpha$  chain of an integrin and the heavy chain or light chain of an immunoglobulin and a recombinant vector for expression of a chimeric protein consisting of the  $\beta$  chain of the integrin and the heavy chain or light chain of the immunoglobulin are transfecting into the cell one after another using different drug resistance

markers. The recombinant vectors can be transfected in any order or simultaneously. It is desirable that the two recombinant vectors to be transfected are vectors for expression of a combination consisting of  $\alpha$  chain • immunoglobulin heavy chain (which means a chimeric protein consisting of an  $\alpha$  chain and the heavy chain of an immunoglobulin; hereinafter this applies) and  $\beta$  chain • immunoglobulin heavy chain, or  $\alpha$  chain • immunoglobulin heavy chain and  $\beta$  chain • immunoglobulin light chain, or  $\alpha$  chain • immunoglobulin light chain and  $\beta$  chain • immunoglobulin heavy chain. Any of these combinations can be adopted, but a combination of recombinant vectors for expression of  $\alpha$  chain • immunoglobulin heavy chain and  $\beta$  chain • immunoglobulin heavy chain is desirable.

[0026] In any transfection method and any combination of vectors, it is important to select a cell which is transfected by the two recombinant vectors and produces a chimera protein consisting of an  $\alpha$  chain and the heavy chain or light chain of an immunoglobulin and a chimera protein consisting of a  $\beta$  chain and the heavy chain or light chain of an immunoglobulin simultaneously almost by the same amounts. This can be achieved by measuring the amounts of the chimeric protein consisting of an  $\alpha$  chain and the heavy chain or light chain of an immunoglobulin and a chimeric protein consisting of a  $\beta$  chain and the heavy chain or light chain of an immunoglobulin produced in the cultured supernatant solution of the cell transfected by the recombinant vectors. For measurement, for example, the transfected cell can be cultured in a medium containing  $^{35}\text{S}$  according to any publicly known method, for labeling the proteins, and the amounts of the chimeric protein consisting of an  $\alpha$  chain and the heavy chain or light chain of an immunoglobulin and a chimeric protein consisting of a  $\beta$  chain and the heavy chain or light chain of an immunoglobulin existing in the cultured supernatant solution can be estimated by immunoprecipitation using an anti- $\alpha$  chain antibody or an anti- $\beta$  chain antibody respectively. As another method, the amounts the chimeric protein consisting of an  $\alpha$  chain and the heavy chain or light chain of an immunoglobulin and a chimeric protein consisting of a  $\beta$  chain and the heavy chain or light chain of an immunoglobulin existing in the cultured supernatant solution can be estimated according to the ELISA method using an anti-human immunoglobulin antibody and an anti- $\alpha$  chain antibody or an anti- $\beta$  chain antibody. Anyway, it is preferable to select a clone which produces almost the same large amounts of the chimeric proteins of the  $\alpha$  and  $\beta$  chains in the culture supernatant, for preparing an integrin-immunoglobulin chimeric protein heterodimer complex. The methods for labeling proteins, the methods of immunoprecipitation and the general methods of ELISA are described in a published book ("Antibody" Harlow, E., and Lane, D. (1988), Cold Spring Harbor Lab. Press, New York), but the methods are not limited to them. Any other method can also be used for detecting chimeric proteins.

[0027] The obtained transfected cell can be cultured according to a general cell culture method, to produce an integrin-immunoglobulin chimeric protein heterodimer complex. It is preferable that the medium contains about 5% of serum of a low immunoglobulin concentration, but any generally known serum-containing medium or a serum-less medium can also be used. After completion of cell culture, the cells and solid matter are removed by such operation as centrifugation, and the culture supernatant containing an integrin-immunoglobulin chimeric protein heterodimer complex is collected.

[0028] It can be estimated that the cultured supernatant solution contains not only the integrin-immunoglobulin chimeric proteins in which the chimeric protein consisting of an  $\alpha$  chain and the heavy chain or light chain of an immunoglobulin and a chimeric protein consisting of a  $\beta$  chain and the heavy chain or light chain of an immunoglobulin form a heterodimer complex, but also the chimeric protein consisting of an  $\alpha$  chain and the heavy chain or light chain of an immunoglobulin and a chimeric protein consisting of a  $\beta$  chain and the heavy chain or light chain of an immunoglobulin which do not form the heterodimer complex. However, since the molecules other than the heterodimer complex cannot be bound to a ligand, the supernatant solution can be used as a reagent for testing the binding to a ligand or cell, or searching for a substance inhibiting the binding between an integrin and a ligand, or for searching for a substance capable of being bound to an integrin, or for measuring the ligand amount of an integrin. These methods of utilization are basically the same as those for using a purified integrin-immunoglobulin chimeric protein heterodimer complex described later.

[0029] An integrin-immunoglobulin chimeric protein heterodimer complex can be purified by an established method using a protein A column chromatography by use of the nature of the immunoglobulin portion. Furthermore, affinity chromatography using an antibody against the  $\alpha$  or  $\beta$  chain can also be used. Moreover, the purification can also be achieved by affinity chromatography with a ligand bound to a carrier. General chromatographic methods can also be used in combination for the purification. If publicly known cases in which integrin molecules are purified by these methods (Pytela, R. et al., *Methods Enzymol.*, 144, 475-489 (1987), Santoro, S. A. et al., *Biochem. Biophys. Res. Comm.*, 153, 217-223 (1988), Charo, I.F. et al., *J. Cell Biol.*, 111, 2795-2800 (1990), Makarewicz, R. et al., *J. Biol. Chem.*, 269, 4005-4011 (1994), Pfaff, M. et al., *Eur. J. Immunol.*, 225, 975-984 (1994), Gulino, D. et al., *Eur. J. Biochem.*, 227, 108-115 (1995), etc.) are applied, the purification of an integrin-immunoglobulin chimeric protein heterodimer complex can be achieved.

[0030] A purified integrin-immunoglobulin chimeric protein heterodimer complex can be identified as a protein showing at least one band under non-reducing condition and at least two bands under reducing condition by SDS-PAGE. It can also be confirmed from it, that the heterodimer is linked by the disulfide bond between immunoglobulin heavy chains. It sometimes occurs that a plurality of bands are detected under reduction, but this is considered to be probably



because intramolecular cleavage of the  $\alpha$  chain has occurred. Especially, with  $\alpha 4$ , this phenomenon is known (Hemler, M. E. et al., J. Biol. Chem., 262, 11478-11485 (1987)). Furthermore, it can be confirmed by the Western blotting method that the respective bands indicate chimeric proteins. As another method, it can be confirmed by said ELISA method combining an anti- $\alpha$  chain antibody, anti- $\beta$  chain antibody and anti-human immunoglobulin antibody, that the obtained molecule is an integrin-immunoglobulin chimeric protein heterodimer complex. That is, the molecule can be identified as a protein molecule with epitopes for all the antibodies. As a further other method, an integrin-immunoglobulin chimeric protein heterodimer complex can also be identified by immunoprecipitation. In this case, if the purified protein is labeled by  $^{35}\text{S}$ , or  $^{125}\text{I}$  or biotin, etc. according to any known method, and immunoprecipitated using an anti- $\alpha$  chain antibody, anti- $\beta$  chain antibody and anti-human immunoglobulin antibody, the same electrophoretic pattern can be obtained in every case. So, it can be confirmed that the integrin-immunoglobulin chimeric protein heterodimer complex has the intended structure. Furthermore, even if a condition to dissociate the integrin complex on a cell membrane such as the coexistence of EDTA or boiling in the presence of SDS is applied, the immunoprecipitation pattern is not changed. So, it can be confirmed that the obtained integrin-immunoglobulin chimeric protein heterodimer complex is a structurally stabilized complex. The methods for confirming an integrin-immunoglobulin chimeric protein heterodimer complex are not limited to those stated above.

[0031] The binding between a prepared integrin-immunoglobulin chimeric protein heterodimer complex and a ligand can be tested as described below. After a ligand and an integrin-immunoglobulin chimeric protein heterodimer complex are brought into contact with each other, to make a mixture, the amount of the integrin-immunoglobulin chimeric protein heterodimer complex bound to the ligand or the amount of the ligand bound to the integrin-immunoglobulin chimeric protein heterodimer complex is measured. The amount of an integrin-immunoglobulin chimeric protein heterodimer complex can be measured by labeling the complex itself by a fluorescent dye or enzyme or radio isotope, etc. The amount of a ligand can also be measured by any similar method. A detection method such as SPA (Amasham) can also be used for the measurement. Furthermore, any reagent which can recognize a complex or ligand labeled by a fluorescent dye, enzyme or radioisotope, etc. can also be used for the measurement. The reagent for recognizing an integrin-immunoglobulin chimeric protein heterodimer complex can, for example, be an anti-human immunoglobulin antibody. In this test, it is preferable to bind the molecule to be detected, to any carrier such as a bead or plate. As a ligand, its entire molecule can be used, but a portion retaining the binding activity to an integrin can also be taken out for use. For example, for integrin  $\alpha 4 \beta 1$  or integrin  $\alpha 2 \beta 1$ , its ligand, fibronectin or collagen or its peptide fragment bound to a carrier can also be used.

[0032] Methods similar to the above can be used to test the binding between an integrin-immunoglobulin chimeric protein heterodimer complex and cells. The amount of the cells bound to a complex can be measured by labeling the cells by a fluorescent dye or radioisotope or using a reagent reacting with the cells, for example, an antibody reacting with a surface antigen. If something like a tissue section is used instead of cells, the amount of the bound integrin-immunoglobulin chimeric protein heterodimer complex is measured by any of the above mentioned methods.

[0033] The methods for examining the binding between an integrin-immunoglobulin chimeric protein heterodimer complex and a ligand or cell described above can be used for obtaining a substance inhibiting the binding between an integrin and a ligand, for example, for obtaining an antibody, polypeptide, peptide or low molecular weight compound. It is preferable to premix a sample and an integrin-immunoglobulin chimeric protein heterodimer complex, and then to measure the amount of the integrin-immunoglobulin chimeric protein heterodimer complex bound to a ligand in any of the above mentioned measuring systems. If the amount of the bound integrin-immunoglobulin chimeric protein heterodimer complex is lowered by adding a certain sample, it can be judged that the sample has inhibitory activity. However, in this system, a substance with metal ion chelating action or a substance with surfactant action, etc. may give a false positive result. The sources of samples used include the following integrin bound substances, peptide fragments of ligands, their derivatives, marketed compounds, etc., but are not limited to them.

[0034] A case where a purified integrin was coated on a plate to search for a peptide to be bound was reported (Healy, J. M. et al., Biochemistry 34, 3948-3955 (1995)). Even if the integrin-immunoglobulin chimeric protein heterodimer complex obtained in the present invention is used, a substance to be bound to an integrin can be similarly searched for. Especially when the chimeric protein heterodimer complex of the present invention is used, the operation to remove the non-specifically bound substances can be effected under more severe conditions. So, the operation can be simplified advantageously. Furthermore, since the complex is not dissociated during operation, a bound substance can be selected more specifically advantageously. Known sources suitable for selecting bound substances include a phage peptide library (e.g., Scott, J. K. and Smith, G. P., Science, 249, 386-390 (1990)) and a DNA oligomer library (e.g., O'Connell, D. et al., Proc. Natl. Acad. Sci. USA, 93, 5883-5887 (1996)), but in the present invention, it is preferable to use the former.

[0035] Furthermore, the method of testing the binding between an integrin-immunoglobulin chimeric protein heterodimer complex and a ligand or cell can also be used as a method for measuring the amount of an integrin ligand in a body fluid or tissue.

[0036] Moreover, the integrin-immunoglobulin chimeric protein heterodimer complexes of the present invention can

also be used as drugs. The present invention has clarified that integrins and other isolated extracellular matrix receptors can be used as platelet substitutes.

[0037] An extracellular matrix receptor preferably used as a platelet substitute is an integrin. The  $\alpha$  chain of the integrin can be  $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 3$ ,  $\alpha 4$ ,  $\alpha 5$ ,  $\alpha 6$ ,  $\alpha 7$ ,  $\alpha 8$ ,  $\alpha 9$ ,  $\alpha v$ ,  $\alpha L$ ,  $\alpha M$ ,  $\alpha X$ ,  $\alpha I b$  or  $\alpha E$ , and among them,  $\alpha 2$  is preferable. The  $\beta$  chain can be  $\beta 1$ ,  $\beta 2$ ,  $\beta 3$ ,  $\beta 4$ ,  $\beta 5$ ,  $\beta 6$ ,  $\beta 7$  or  $\beta 8$ , and among them,  $\beta 1$  is preferable. Integrin  $\alpha 2 \beta 1$  is more preferable. The receptor source for isolation can be a tissue or cell expressing an extracellular matrix receptor, or a dissolved membrane fraction of a receptor expressing cell prepared by gene manipulation, etc. It is more preferable to design for obtaining a soluble protein by modifying a receptor gene by gene recombination, and to use the cultured supernatant solution of the cells capable of producing it, as a source. Furthermore in the design of the soluble protein, it is preferable that the functional structure of the extracellular matrix receptor is retained. For example, it is desirable to use an integrin-immunoglobulin chimeric protein heterodimer complex obtained by modifying the heterodimer structure of an integrin to allow its  $\alpha$  and  $\beta$  chains to be covalently associated with each other. As the integrin-immunoglobulin chimeric protein heterodimer complex, it is preferable that the  $\alpha$  chain of the integrin is  $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 3$ ,  $\alpha 4$ ,  $\alpha 5$ ,  $\alpha 6$ ,  $\alpha 7$ ,  $\alpha 8$ ,  $\alpha 9$ ,  $\alpha v$ ,  $\alpha L$ ,  $\alpha M$ ,  $\alpha X$ ,  $\alpha I b$  or  $\alpha E$ , and among them,  $\alpha 2$  is more preferable. Furthermore, it is preferable that the  $\beta$  chain is  $\beta 1$ ,  $\beta 2$ ,  $\beta 3$ ,  $\beta 4$ ,  $\beta 5$ ,  $\beta 6$ ,  $\beta 7$  or  $\beta 8$ , and among them,  $\beta 1$  is more preferable. It is further more preferable that the  $\alpha$  chain is  $\alpha 2$  and that the  $\beta$  chain is  $\beta 1$ . The platelet substitute of the present invention is described below mainly in reference to a typical extracellular matrix receptor, integrin  $\alpha 2 \beta 1$ -immunoglobulin chimeric protein heterodimer complex, but the present invention is not limited thereto or thereby.

[0038] To confirm the applicability of a purified integrin-immunoglobulin chimeric protein heterodimer complex as a drug, the purified protein itself is used for examining its pharmacological activity. For obtaining higher capability of being bound to an extracellular matrix, it is more preferable to use an integrin-immunoglobulin chimeric protein heterodimer complex bound to a carrier such as a lipid or protein polymer, etc., but the present invention is not limited to this method.

[0039] For use as a platelet substitute, it is preferable to bind an integrin  $\alpha 2 \beta 1$ -immunoglobulin chimeric protein heterodimer complex to a liposome covalently according to the method stated in a report (Martin, F. J. et al., Biochemistry, 20, 4229 (1981)). The carrier can also be any other drug carrier than a liposome as far as its use for drugs is permitted. If a liposome is used as the carrier, the liposome is prepared according to the composition and method stated in a published book "Preparation and Experiments of Liposomes (in Japanese)", Oku, N. (1994), Hirokawa Shoten, but a preferable method is such that the epitope bound to the extracellular matrix of an integrin  $\alpha 2 \beta 1$ -immunoglobulin chimeric protein heterodimer complex is exposed outside the liposome membrane.

[0040] For confirming that an integrin  $\alpha 2 \beta 1$ -immunoglobulin chimeric protein heterodimer complex is bound on the prepared liposome carrier, a flow cytometer is used. The reagents which can be used for recognizing the integrin  $\alpha 2 \beta 1$ -immunoglobulin chimeric protein heterodimer complex include an anti-integrin  $\alpha 2$  antibody, anti-integrin  $\beta 1$  antibody, anti-human immunoglobulin antibody, etc. If the antibody used is fluorescently labeled, it can be used for determination directly, but if it is not fluorescently labeled, a secondary antibody which recognizes the immunoglobulin class of the animal species used for preparing the antibody is used as a fluorescent label. As a further other confirmation method, the integrin  $\alpha 2 \beta 1$ -immunoglobulin chimeric protein heterodimer complex itself can be labeled by an enzyme or radioisotope, etc., for confirmation in proper combination with a color dye or radioactivity measuring instrument, etc.

[0041] To examine the extracellular matrix binding capability using an integrin  $\alpha 2 \beta 1$ -immunoglobulin chimeric protein heterodimer complex liposome, it is preferable to suspend the integrin  $\alpha 2 \beta 1$ -immunoglobulin chimeric protein heterodimer complex liposome into a buffer with a physiological cation concentration or plasma. The buffer with a physiological cation concentration refers to a buffer containing at least cations such as Mg ions or Ca ions and adjusted to about neutrality. The plasma is prepared by processing the blood collected in the presence of an anticoagulant, according to a general plasma preparation method. As the anticoagulant, for example, heparin or EDTA solution can be added by sufficient units. A marketed normal plasma, coagulation factor deficient plasma or serum, etc. can also be used. However, if the anticoagulant used lowers the cation concentration, cations are added to achieve a physiological concentration later. Then, the integrin  $\alpha 2 \beta 1$ -immunoglobulin chimeric protein heterodimer complex liposome is mixed with an extracellular matrix or its fragment coated on a carrier for a certain time, to judge whether binding takes place. It is preferable that the coating of the extracellular matrix or its fragment as a solid phase is achieved by using a plastic plate, etc., but marketed beads for coating an extracellular matrix as a solid phase, etc. can also be used. When a collagen is used as the extracellular matrix, any animal species and type can be used. The binding reaction between an integrin  $\alpha 2 \beta 1$ -immunoglobulin chimeric protein heterodimer complex liposome and an extracellular matrix is effected according to a general method adopted for observing the adherence reaction of platelets. In many cases, they are allowed to stand mainly in a static system for a certain time, to induce binding to the matrix, but it is preferable to apply a shaking or shear stress, etc.

[0042] The integrin  $\alpha 2 \beta 1$ -immunoglobulin chimeric protein heterodimer complex liposome is bound to an extracellular matrix under the conditions as described above, and the amount of binding is measured by applying the above mentioned ELISA method using an anti-human immunoglobulin antibody. For more accurate determination, it is desirable to



immobilize the liposome bound to the matrix by 1% glutaraldehyde, etc. As another method than the ELISA method, for example, if a radio-labeled lipid is incorporated into the liposome beforehand, the amount of the liposome bound to the extracellular matrix can be obtained as radioactivity. Furthermore, to qualitatively judge the binding and covering degree to the extracellular matrix, a labeled antibody for recognizing the integrin  $\alpha 2\beta 1$ -immunoglobulin chimeric protein heterodimer complex on the bound liposome can be combined with a color dye, etc., to dye the portions where the liposome is bound. It is more preferable that the generally used tissue antibody dyeing method is used to use a peroxidase labeled antibody against the integrin  $\alpha 2\beta 1$ -immunoglobulin chimeric protein heterodimer complex and diaminobenzidine in combination, but the measuring method is not limited to it. As a further other method, the area covering the extracellular matrix can be obtained as a covering rate using an image processing analyzer.

[0043] Methods for examining the hemostasis of platelets include testing the adhering capability of platelets to the extracellular matrix and the agglutination capability induced by a collagen ("Handbook on the Examination of Blood Coagulation (in Japanese)", p. 65-78, Fukutake, M. and Fujimaki, M. (1987), Uchudo Yagi Shoten, Santro, S.A., Cell, 46, 913-920 (1986), Lethagen, S. and Rugarrn, P., Thrombo Haemost., 67, 185-186 (1982)). Especially the adhering capability of platelets to the extracellular matrix is an indicator of primary hemostasis. The adhering capability is evaluated by using blood as it is, or platelet rich plasma or platelets washed by a buffer with physiological ions. Therefore, whether or not the integrin  $\alpha 2\beta 1$ -immunoglobulin chimeric protein heterodimer complex liposome obtained in the present invention can be a functional substitute of platelets can be judged in reference its binding capability and the level of the binding capability to the extracellular matrix in the existence of plasma components or at a physiological ion concentration.

[0044] If the binding capability of the integrin  $\alpha 2\beta 1$ -immunoglobulin chimeric protein heterodimer complex liposome obtained in the present invention to the extracellular matrix in the presence of the plasma components is strong, it suggests that the liposome can be a platelet substitute. Therefore, it can be used as a therapeutic or preventive agent against the congenital and acquired bleeding tendency due to platelet abnormality, and also widely as a platelet transfusion substitute.

[0045] Similarly the integrin  $\alpha 2\beta 1$ -immunoglobulin chimeric protein heterodimer complex liposome obtained in the present invention can be a therapeutic or preventive agent for conditions of diseases where vascular endothelial cell disorder is a problem. For example, it was reported that in the prognosis of PTCA (percutaneous coronary restenosis), the excessive accumulation of platelets on the extracellular matrix exposed by balloon catheter treatment triggers restenosis (Liu, M.W. et al., Circulation, 79, 1374-1378 (1989)). In Example 22, the effect of the integrin  $\alpha 2\beta 1$ -immunoglobulin chimeric protein heterodimer complex liposome to cover the extracellular matrix was confirmed, and this effect can reduce the excessive accumulation of platelets to allow use also as a restenosis preventive. Furthermore, if the integrin  $\alpha 2\beta 1$ -immunoglobulin chimeric protein heterodimer complex liposome is labeled by a medically allowable method, it can be used for monitoring the region of the extracellular matrix exposed by vascular endothelial cell injury, and furthermore, if a drug is enclosed in the liposome, it can also be applied to the targeting therapy for a local injured region.

[0046] When any integrin  $\alpha 2\beta 1$ -immunoglobulin chimeric protein heterodimer complex liposome stated in the present invention is used as a platelet substitute, the administration paths include infusion, intravenous administration, etc., and it is usually used by being suspended in any physiologically suitable solution such as a salt solution or plasma, etc. It can be used alone or also in combination with another chimeric protein heterodimer complex with an extracellular matrix receptor or its immunoglobulin. It can also be used together with another drug containing total platelets. The dose is properly selected to suit the symptom, age, body weight, etc. and can be 0.1 mg to 10 g per day as the amount of the protein for an adult, being able to be administered at a time or in several times. It can also be mixed with a pharmaceutically allowed carrier or excipient, etc., to be applied locally to the injured region as an externally applied drug such as an ointment, liniment or plaster. In this case, the externally applied drug is prepared to be 1 ng/cm<sup>2</sup> to 1 mg/cm<sup>2</sup> as the amount of the protein per one time of coating.

#### Examples

[0047] To describe the present invention in more detail, examples are given below. The general methods of recombinant DNA experiments conformed to those stated in a published book ("Antibody", Harlow, E. and Lane, D. (1988), Cold Spring Harbor Lab. Press, New York).

#### Example 1

##### Construction of human IgG<sub>1</sub> heavy chain expression vector

[0048] As human IgG<sub>1</sub> genome gene, a clone identical with reported base sequence information (Ellison, J. W. et al., Nucleic Acids Res., 10, 4071-4079 (1982)) was acquired from a human genomic library (CLONTECH) using a hybridization cDNA probe based on the sequence information. This was used as the template DNA for PCR. As primers for

amplifying the DNA fragment containing the hinge region (H) and the constant region portions (CH2 and CH3) of human IgG<sub>1</sub> gene, a DNA oligomer shown in sequence No. 4 of the sequence table (hereinafter a sequence No. of the sequence table is simply called a sequence No.) with BamH I restriction site and a DNA oligomer shown in sequence No. 5 with Xba I restriction site were synthesized.

5' - GCGGATCCCGAGCTGCTGGAAGCAGGCTCAG - 3' (Sequence No. 4)

5' - CCTCTAGACGGCCCTCGCACTCATTTA - 3' (Sequence No. 5)

[0049] The template DNA, primers, dNTPs (an equimolar mixture of dATP, dCTP, dGTP and dTTT) and Taq polymerase (Takara) were mixed in a PCR buffer (100 mM Tris-HCl, 500 mM KCl, 15 mM MgCl<sub>2</sub>, 0.01% gelatin, pH 8.3), and in a thermal cycler (Perkin Elmer Cetus), the mixture was treated at 94 °C for 1 minute for DNA denaturation, at 58°C for 2 minutes for annealing the primers and at 72°C for 3 minutes for elongating the primers. This treatment was performed 30 cycles. The amplified DNA was digested by restriction enzymes BamH I and Xba I, and the DNA fragment was purified by 1% agarose gel according to a general method ("Antibody", Harlow, E. and Lane, D. (1988), Cold Spring Harbor Lab. Press, New York). It was linked, using a T4DNA ligase, with a large DNA fragment of pBluescriptSK(+) (STRATAGENE) purified and digested by restriction enzymes BamH I and Xba I. The plasmid DNA was used to transform Escherichia coli (JM109), and the transformant was selected, to obtain a plasmid DNA (IgG<sub>1</sub> Bluescript). Then, expression vector pcDL-SR α296 was digested by restriction enzyme BamH I, and blunted at the termini by T4DNA polymerase treatment, and a Not I linker was linked. The large DNA fragment obtained by digesting it by restriction enzymes Not I and Xho I and the small DNA fragment obtained by digesting IgG<sub>1</sub> Bluescript by restriction enzymes Not I and Xho I were purified according to a general method, and linked by T4DNA ligase. It was transformed into Escherichia coli (HB101), and the transformant was selected, to obtain a plasmid DNA. Hereinafter this plasmid (IgG<sub>1</sub> SRα) is called human IgG<sub>1</sub> expression vector. In the following examples, since the basic protocol of gene manipulation is the same as above, the description will be simplified.

## Example 2

### Construction of integrin α4 - IgG heavy chain chimeric protein expression vector

[0050] The DNA fragment coding for the extracellular portion of integrin α4 was obtained by cloning based on reported cDNA sequence information (Takada, Y. et al., EMBO J., 8, 1361-1368 (1989)). The restriction site EcoR I of 1801-base-position of sequence No. 1, the restriction site Stu I of 112-base-position and the restriction site BamH I of 2949-base-position were used for linking the region from the N terminus translation initiation site to Stu I cut site as α4-1, the region from Stu I cut site to EcoR I cut site as α4-2, and the region from EcoR I detached site to BamH I detached site as α4-3. The detailed methods are described below.

[0051] The portion coding for α4-1 was designed to be cloned by linking the DNA oligomers of sequence Nos. 6 to 9, and the DNA oligomers shown in sequence Nos. 6 to 9 were synthesized. For the sequence Nos. 6 and 7, restriction site Xba I was added on the side to code for the N terminus, for linking to a vector. Furthermore, compared with the known sequence information, the bases at the 60-, 63- and 64-positions were substituted from C to T, C to A and C to G respectively, and the bases at the 112- and 114-positions were substituted from C to A and C to G respectively. Because of substitution at the 112- and 114-position, restriction site Stu I was inserted on the side to code for the N terminus of sequence Nos. 8 and 9. The 5' termini of the synthesized oligomers were phosphorylated and annealed, and were linked using T4DNA ligase. After completion of linking, restriction enzymes Xba I and Stu I were used for cutting, and electrophoresis was effected by 5% agarose (NuSieve GTGagarose, FMC) gel. The intended DNA fragment (α4-1) of about 120 bp was cut out and purified.

5' - CTAGACCACCATGTTCCCCACCGAGAGCCCATGGCTTGGCAAGCGAGCGCGGAACCCGGCCCCCGGA  
GCTGCA-3' (Sequence No. 6)

5' - GCTTCGGGGCCCCGGTTCGGCCTCGCTTCCCAAGCCATGCCCTCTCGGTGGGAAACATGGTGGT-3' (Sequence No. 7)

5' - CTCCGGCAGACGGTCATGCTGTTGCTGTGCCTGGGGTCCCGACCGGCAGC-3' (Sequence No. 8)

5' - CCTCCGGTCCGGACCCCCAGGCACAGCAACAGCATCACCGTCTCCCGAGTCCA-3' (Sequence No. 9)

[0052] Then, the RNA of human osteosarcoma cell line MG63 (ATCC CRL 1427) as an integrin  $\alpha 4$  expressing cell was separated, and PolyA(+) RNA was purified using oligo dT cellulose column (NEB). Based on it, a single stranded cDNA was synthesized using a reverse transcriptase (GIBCO), and used as the template for PCR. As primers for amplifying  $\alpha 4$ -2 and  $\alpha 4$ -3 DNAs, four DNA oligomers of sequence Nos. 10 to 13 with Pst I and Stu I restriction sites inserted (sequence No. 10) or BamH I restriction site inserted (sequence No. 13) were synthesized:

5' - CACTCCAGGCAGGCTTACAACGTCCACACTGACAGC-3' (Sequence No. 10)

5' - CCAGAAACCTCTAAATCAGCAG-3' (Sequence No. 11)

5' - GCATTATCGCGAAAGATCTGC-3' (Sequence No. 12)

5' - CGGCATCCGTGAAATAACGTTTGGGTCTT-3' (Sequence No. 13)

[0053] The template cDNA, primers, dNTPs and Taq polymerase were mixed in a PCR buffer, and the mixture was treated by a thermal cycler at 94°C for 1 minute for DNA denaturation, at 58°C for 2 minutes for annealing the primers and at 72°C for 3 minutes for elongating the primers. This treatment was performed 30 cycles. The amplified DNA fragments of  $\alpha 4$ -2 and  $\alpha 4$ -3 were digested by Pst I and EcoR I respectively, or EcoR I and BamH I and sub-cloned into pBluescriptKS(+) (STRATAGENE), to prepare plasmid DNAs (hereinafter called  $\alpha 4$ -2 Bluescript and  $\alpha 4$ -3 Bluescript). Then, upstream of the  $\alpha 4$ -2 Bluescript,  $\alpha 4$ -1 was linked using Xba I and Stu I restriction sites, to prepare a plasmid DNA (hereinafter called  $\alpha 4$ -1-2 Bluescript).

[0054] The  $\alpha 4$ -1-2 Bluescript was digested by restriction enzyme Not I, and blunted at the termini by T4DNA polymerase treatment, being digested by restriction enzyme EcoR I, to prepare a small DNA fragment. The  $\alpha 4$ -3 Bluescript was digested by restriction enzymes EcoR I and BamH I, to prepare a small DNA fragment. The two small DNA fragments were simultaneously linked to a large DNA fragment obtained by digesting IgG<sub>1</sub>SR $\alpha$  by restriction enzymes EcoR V and BamH I, to obtain a plasmid DNA. The obtained base sequence coding for integrin  $\alpha 4$  - IgG heavy chain chimeric protein is shown as sequence No. 1. The plasmid (integrin  $\alpha 4$  - IgGSR $\alpha$ ) is hereinafter called integrin  $\alpha 4$  - IgG heavy chain chimeric protein expression vector.

## Example 3

Construction of  $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein expression vector

5 [0055] The RNA of human fibroblast cell line MRC5 (ATCC CCL 171) as an integrin  $\beta 1$  expressing cell was separated, and oligo dT cellulose column was used to purify PolyA (+)RNA. Based on it, a single stranded cDNA was synthesized using a reverse transcriptase, and used as the template for PCR. As primers, two DNA oligomers of sequence Nos. 14 and 15 with BamH I site (sequence No. 15) inserted on the side coding for C terminus were synthesized according to the sequence information (Cott Argraves, W. et al., J. Cell Biol., 105, 1183-1190 (1987)).

10

5' - GCGGAAAAGATGAATTTACAAC-3' (Sequence No. 14)

5' - GTGGGATCCTCTGGACCACTGGGACAC-3' (Sequence No. 15)

15

[0056] The template cDNA, primers, dNTPs and Taq polymerase were mixed in a PCR buffer, and treated by a thermal cycler at 94°C for 1 minute for DNA denaturation, at 57°C for 2 minutes for annealing the primers and at 72°C for 3 minutes for elongating the primers. This treatment was performed 30 cycles. The amplified DNA was blunted at the terminus by T4DNA polymerase treatment, and digested by restriction enzyme BamH I. Then, the DNA fragment was purified. Subsequently, the DNA fragment obtained in PCR before was sub-cloned at the Sma I and BamH I sites of pBluescriptKS(+). A small DNA fragment purified by digesting it by restriction enzymes EcoR I and BamH I was inserted into a large DNA fragment of IgG<sub>1</sub>SR $\alpha$  treated by restriction enzymes EcoR I and BamH I, to obtain a plasmid DNA.

25 The obtained base sequence coding for  $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein is shown in sequence No. 2. The plasmid (integrin  $\beta 1 \cdot \text{IgGSR}\alpha$ ) is hereinafter called integrin  $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein expression vector.

## Example 4

30 Transfection of  $\alpha 4 \cdot \text{IgG}$  heavy chain chimeric protein expression vector and  $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein expression vector into animal cells, and their expression

[0057] Integrin  $\beta 1 \cdot \text{IgGSR}\alpha$  as  $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein expression vector and pSV2dhfr (BRL) were mixed at a ratio of 10 : 1, and the mixture and lipofectin reagent (GIBCO BRL) were gently mixed and allowed to stand at room temperature for 15 minutes. The mixture was added dropwise to dihydrotolic acid reductase deficient CHO cells (ATCC CRL 9096). After 18 hours of dropwise addition, the mixture was cultured in a medium (10%FBS (GIBCO), nucleic acid-containing  $\alpha$ MEM medium (GIBCO BRL)) for about 2 days, and the cells were dispersed by trypsin-EDTA treatment. The cells were suspended in a first selective medium (10% FBS-containing nucleic acid-free  $\alpha$ MEM medium (GIBCO BRL)), and the suspension was disseminated into a 96-well plate (CORNING), for selective culture for about 10 days. Then, the amount of integrin  $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein produced in the culture supernatant was determined according to the ELISA method (described later), and the clone showing the highest production was stabilized by cloning according to the limiting dilution method.

[0058] Then, into the stabilized integrin  $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein producing CHO cells, the integrin  $\alpha 4 \cdot \text{IgG}$  heavy chain chimeric protein expression vector was transfected according to the lipofectin method as described before. That is, integrin  $\alpha 4 \cdot \text{IgGSR}\alpha$  and pSV2neo (BRL) were mixed at 10 : 1, and the mixture was mixed with lipofectin reagent. The mixture was added dropwise into the cells. After 18 hours of dropwise addition, the mixture was cultured in the said first selective medium for about 2 hours, and the cells were dispersed by trypsin-EDTA treatment. The cells were suspended in a second selective medium (nucleic acid-free  $\alpha$ MEM medium (GIBCO BRL) containing 10% FBS (GIBCO) and 1 mg/ml neomycin (GIBCO)), and on a 96-well plate (CORNING), resistant cells were selectively cultured for about 10 days. The amount of integrin  $\alpha 4 \cdot \text{IgG}$  heavy chain chimeric protein and the amount of integrin  $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein produced in the culture supernatant were determined according to the ELISA method (described later), and a clone which produced both the chimeric proteins by almost the same amounts was picked up. The clone was cloned twice according to the limiting dilution method, to be stabilized as a clone capable of producing  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex.

55

## Example 5

Determination of produced integrin  $\alpha 4 \cdot$  IgG heavy chain chimeric protein and integrin  $\beta 1 \cdot$  IgG heavy chain chimeric protein by the ELISA method

- 5 [0059] Fifty microliter per well of anti-human integrin  $\alpha 4$  antibody (Becton & Dickinson, Clone L25.3) or anti-human integrin  $\beta 1$  antibody (Coulter, Clone 4B4) (12  $\mu$ g/ml each) was put into a 96-well immunoplate (NUNC) and allowed to stand at 4°C for 16 hours. Then, each well was washed by Dulbecco's phosphate buffered saline (Nissui Seiyaku, not containing Ca or Mg ions, hereinafter called PBS(-)) twice, and non-specific reaction was blocked by PBS(-) containing 25% Block Ace (Snow Brand Milk Products Co., Ltd.). After blocking, the culture supernatant of CHO cells grown in selective medium was properly diluted, and reacted with the coated antibody at room temperature for 1 hour. After the reaction, the surface of the plate was washed with 0.02% Tween-containing PBS(-) (hereinafter called T-PBS) twice. It was then caused to react with biotinized anti-human IgG antibody (Vector) for 1 hour, and the reaction mixture was washed with T-PBS twice, and in succession caused to react with avidin-horseradish peroxidase (Sigma) for 1 hour. The reaction mixture was washed with PBS(-) twice. The PBS(-) was perfectly aspirated, and or thophenylenediamine was used as a substrate for color development. The absorbance at 490 nm were measured using a microplate reader (Bio-rad NOVAPATH), and the clone showing a high absorbance value was selected.

## Example 6

20

Purification of  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex

(1) Culture of CHO cells and preparation of cultured supernatant solution

- 25 [0060] The CHO cells highly capable of producing the  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex were cultured in nucleic acid-free  $\alpha$ MEM medium containing 5% FBS (Ultra-low IgG grade, GIBCO) (hereinafter called  $\alpha$ MEM(-) medium, GIBCO BRL) for one day, to reach semiconfluent, and they were cultured in  $\alpha$ MEM(-) medium containing 1% FBS (Ultra-low IgG grade) for 3 days, and the culture supernatant was collected. It was concentrated to 1/10 volume by ultrafiltration using Prep-scale (Millipore), and 1M Hepes solution (pH 8.0) was added to achieve a final concentration of 5 mM, for preparing a starting solution for further purification.

(2) Protein A column chromatography

- 35 [0061] The starting solution for further purification was passed through Prosep Guard column (bioPROCESSING), and applied to Prosep A column (bioPROCESSING). After completion of application, it was washed with 10 times the column volume of PBS(-), and the proteins were eluted at a pH 6-3 gradient of 0.1M citrate buffer solutions. The peak fraction eluted at pH 3 was collected, and 1M Tris-HCl solution (pH 8.5) was added by 0.1 volume for neutralization. The solution was dialyzed against PBS(-).

40 (3) Affinity column chromatography

- 45 [0062] FMP activated Cellulofine (Seikagaku Kogyo) was equilibrated by a coupling buffer (50 mM  $\text{Na}_2\text{CO}_3$ - $\text{NaHCO}_3$  pH 8.5), and a peptide showing sequence No. 3 (hereinafter called CS-1 peptide) synthesized by a peptide synthesizer was added. The mixture was inverted and mixed at 4 °C for 16 hours.

Cys Leu His Gly Pro Glu Ile Leu Asp Val Pro Ser Thr (Sequence No. 3)

- 50 After completion of mixing, the mixture was washed with the coupling buffer, and a blocking buffer (0.1 mM monoethanolamine, 50 mM Tris-HCl, pH 8.0) was added. The mixture was inverted and mixed further at room temperature for 6 hours. Then, the mixture was sufficiently washed with TBS solution (150 mM NaCl, 20 mM Tris-HCl, 1 mM  $\text{MnCl}_2$ , pH 7.5), to prepare CS-1 peptide bound Cellulofine column. To the column the starting solution for further purification was applied and allowed to stand at room temperature for 3 hours, and washed with 10 times the column volume of a washing buffer (1M NaCl, 0.1% Triton, 20 mM Tris-HCl, 1 mM  $\text{MnCl}_2$ , pH 7.5) and the same volume of the TBS solution. After completion of washing, an elution buffer (10 mM EDTA, 150 mM NaCl, 20 mM Tris-HCl, pH 7.5) was used, to elute the proteins bound to the CS-1 column. The eluate was collected and dialyzed against PBS(-).



## (4) SDS-PAGE

[0063] The eluted fractions of (3) were subjected to SDS-PAGE under non-reducing or reducing condition using 6.0 or 7.0% acrylamide gel, and the gel was stained with Coomassie-blue. As a result, under non-reducing condition, two bands considered to be attributable to the  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex and its polymer were observed. Under reducing condition, two bands (170 kDa and 135 kDa) considered to be attributable to the integrin  $\alpha 4 \cdot \text{IgG}$  heavy chain chimeric protein and the integrin  $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein and two bands (80 kDa and 90 kDa) considered to be attributable to the intramolecular cleavage of the integrin  $\alpha 4 \cdot \text{IgG}$  heavy chain chimeric protein (Hentler, M. E. et al., J. Biol. Chem., 262, 11478-11485 (1987)) were observed. These results suggest that the eluted protein of (3) has a molecular structure considered to be  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex, and that the molecules constituting the heterodimer are linked by a disulfide bond between the IgG heavy chains.

## Example 7

Identification of  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex, and examination of its structural stability

## (1) Immunoprecipitation using anti-integrin antibodies and influence of a cationic chelating agent

[0064] The basic method conformed to a published book ("Antibodies", Harlow, E. et al., (1988), Cold Spring Harbor Lab. Press, New York). That is, the eluted protein of Example 6 (3) considered to be  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex was  $^{125}\text{I}$ -labeled using the lactoperoxidase method. Then, Affigel-10 (Bio-rad) was washed with 0.1 M Hepes solution (pH 8.0), and normal murine IgG, anti-human integrin  $\alpha 4$  antibody (clone 11C2B) and anti-human integrin  $\beta 1$  antibody (clone 4B4) were added. Reaction was effected at 4°C for 16 hours to cause covalent bonding, to prepare normal murine IgG beads and the respective antibody beads. Then, the  $^{125}\text{I}$  labeled  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex and normal murine IgG beads were inverted and mixed at 4 °C for 4 hours for preclearing, and the mixture and the antibody beads were inverted and mixed at 4°C for 16 hours. After completion of mixing, the beads were washed with a washing buffer (200 mM Tris-HCl, 0.5 M NaCl, 0.1% NP-40, 1 mM  $\text{MgCl}_2$  or 10 mM EDTA, pH 8.0) three times. After completion of washing, a sample buffer for electrophoresis was added to the beads for treatment at 100 °C for 5 minutes, and the mixture was centrifuged. The supernatant solution was analysed by electrophoresis under reducing condition. After completion of electrophoresis, the gel was dried by a gel dryer, and the protein was detected by autoradiography.

[0065] As a result of immunoprecipitation in the presence of 1 mM  $\text{MgCl}_2$ , from the beads of both the anti-human integrin  $\alpha 4$  antibody and the anti-human integrin  $\beta 1$  antibody, the same precipitation patterns expected from the structure of  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex were obtained. Thus, the protein obtained in (3) of Example 6 was identified as  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex.

[0066] On the other hand, the immunoprecipitation pattern obtained by using the anti-integrin  $\beta 1$  antibody beads in the presence of 10 mM EDTA was the same as that in the presence of 1 mM  $\text{MgCl}_2$ , to clarify that the association between integrin  $\alpha 4 \cdot \text{IgG}$  heavy chain chimeric protein and integrin  $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein does not depend on cations. The above results suggest that the eluted protein obtained in (3) of Example 6 was certain  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex, and if the result of (4) of Example 6 is also taken into account, it is strongly suggested that the association between both the proteins is stable association through a disulfide bond existing the IgG heavy chains.

(2) Examination on the structural stability of  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex by sequential immunoprecipitation

[0067] According to (1),  $^{125}\text{I}$  labeled  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex was caused to react with normal murine IgG beads, anti-human integrin  $\alpha 4$  antibody (11C2B) beads or anti-human integrin  $\beta 1$  antibody (4B4) beads at 4°C for 4 hours, and the reaction mixture was washed. After washing, the reaction mixture was boiled at 100°C for 5 minutes in the presence of 2% SDS and centrifuged, and the supernatant (primary immunoprecipitation sample) was diluted to 10 times by 1% BSA-containing PBS, and was again reacted with the anti-integrin  $\beta 1$  antibody beads and the anti-integrin  $\alpha 4$  antibody beads at 4°C for 16 hours. After completion of reaction, the beads were washed, and a sample buffer for electrophoresis was added. The mixture was treated at 100°C for 5 minutes and centrifuged. The supernatant solution (secondary immunoprecipitation sample) was analyzed by SDS-PAGE/autoradiography.



[0068] As a result, the electrophoretic pattern obtained by the primary immunoprecipitation was also similarly observed in the secondary immunoprecipitation. This result suggests that the association between the  $\alpha 4 \cdot$  IgG heavy chain chimeric protein and the  $\beta 1 \cdot$  IgG heavy chain chimeric protein in the  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex is not dissociated either by boiling in the presence of 2% SDS, and strongly supports that the complex has a stable heterodimer structure based on a disulfide bond.

#### Example 8

##### Binding of $\alpha 4 \cdot$ IgG heavy chain- $\beta 1 \cdot$ IgG heavy chain chimeric protein heterodimer complex to VCAM-1

[0069] It was examined that the  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex produced by CHO cells can be bound to the ligand of integrin  $\alpha 4 \beta 1$  by using the cells expressing VCAM-1. Human normal umbilical intravenous endothelial cells were cultured with IL-1 3U/ml for 16 hours, to prepare VCAM-1 expressing cells. The cells were treated by 1 mM EDTA at 37°C for 15 minutes, for dispersion as single cells. The cells ( $2 \times 10^5$  cells per sample tube) were cultured with the supernatant of the CHO cells producing  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex for 30 minutes in the presence of 1 mM (final concentration)  $\text{MnCl}_2$  or 3 mM (final concentration) EDTA. After completion of reaction, the cells were washed twice by centrifugation at 1200 rpm at room temperature for 5 minutes using a buffer for binding assay (24 mM Tris-HCl, 10 mM Hepes, 150 mM NaCl, 1 mM  $\text{MnCl}_2$  or 1 mM EDTA, 1% BSA, 2 mM glucose, pH 7.4). After washing, FITC labeled-anti-human IgG antibody (Cappel) was added, and incubated at room temperature for 20 minutes. The cells were washed by the same buffer, and the chimeric proteins bound to the cells were determined by a flow cytometer (ELITE, Coulter).

[0070] The results are shown in Fig. 1. It was observed that the fluorescence intensity showing the binding of  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex increased by culturing the VCAM-1 expression cells with the supernatant containing  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex. The binding was inhibited by adding anti-human integrin antibodies (anti- $\alpha$  antibody: clone L25.3, 10  $\mu\text{M}/\text{ml}$  + anti- $\beta 1$  antibody: clone 4B4, 10  $\mu\text{M}/\text{ml}$ ) or 3 mM EDTA. This result suggests that the  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex can be bound to VCAM-1 like the integrin  $\alpha 4 \beta 1$  existing on the surfaces of cell membranes, and furthermore that the binding is  $\alpha 4 \beta 1$ -specific and retains a feature of the binding that it is dependent on cations.

#### Example 9

##### Binding of $\alpha 4 \cdot$ IgG heavy chain- $\beta 1 \cdot$ IgG heavy chain chimeric protein heterodimer complex to a peptide fragment of fibronectin

[0071] The capability of  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex to be bound to the peptide fragment (sequence No. 3) of the other ligand, fibronectin was also examined.

[0072] At first, according to the said report (Humphries, M. J. et al., J. Biol. Chem., 262, 6886-6892 (1987)), the peptide fragment of sequence No. 3 (CS-1 peptide) was bound to rabbit IgG (Sigma), to prepare CS-1-IgG. The CS-1-IgG was diluted by PBS(-), and put in a 96-well immunoplate (NUNC) by 100  $\mu\text{l}/\text{well}$ , and allowed to stand at 4°C for 16 hours, to be formed as a solid phase on the plate.

[0073] After completion of standing, the surface of the plate was washed with PBS(-) twice and treated with denatured 1% BSA(heat-natured at 80°C for 10 minutes)-PBS solution (300  $\mu\text{l}/\text{well}$ ) at 4°C for 3 hours to block the nonspecific reaction. Then, the solid phase CS-1-IgG and the CHO culture supernatant (100  $\mu\text{l}$ ) containing  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex were reacted with each other at 30°C for 3 hours. The non-bound  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex was removed by washing with 0.1% BSA-containing TBS buffer (150 mM NaCl, 25 mM Tris-HCl, 1 mM  $\text{MnCl}_2$ , pH 7.4) twice, and the bound  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex was detected by biotin labeled anti-human IgG antibody (Vector) as the primary antibody and avidin labeled horseradish peroxidase (Sigma) as the secondary antibody. The surface of the plate was washed with the TBS buffer. Orthophenylenediamine was added as a substrate to it for color development, and the absorbance at 490 nm were measured.

[0074] The results are shown in Fig. 2. The reaction with  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex showed a rise in the absorbance indicating the binding to CS-1 peptide. The binding was almost perfectly inhibited by the presence of anti-integrin  $\alpha A$  antibody (clone L25.3), anti-integrin  $\beta 1$  antibody (clone 4B4) or 5 mM EDTA. Therefore, it was clarified that  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex can be bound also to the CS-1 peptide which is a peptide fragment of fibronectin, and that a feature of binding that it depends on cations is retained.

## Example 10

Evaluation of an inhibitory peptide by using a system for determining the binding of  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex to a peptide fragment of fibronectin

[0075] In the binding determination system of Example 9, the effects of three peptides, i.e., sequence No. 16 (hereinafter called GPEILDVPST), 17 (hereinafter called GPEILEVPST) and 18 (hereinafter called GRGDSP) were examined.

Gly Pro Glu Ile Leu Asp Val Pro Ser Thr (Sequence No. 16)

Gly Pro Glu Ile Leu Glu Val Pro Ser Thr (Sequence No. 17)

Gly Arg Gly Asp Ser Pro (Sequence No. 18)

[0076] The all peptides were synthesized by a peptide synthesizer. The peptide and 100  $\mu$ l of CHO cultured supernatant solution containing  $\alpha 4 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex were mixed at room temperature for 20 minutes, and the binding to CS-1-IgG was determined according to the method in Example 9. The results are shown in Fig. 3. GPEILDVPST showed temperature-dependent inhibitory activity in a range of 0.1 to 10  $\mu$ g/ml, but GPEILEVPST and GRGDSP did not show any inhibition of the binding. These results show that the binding determination system in Example 9 allows to detect the inhibiting effect of the peptide (GPEILDVPST) inhibiting the binding between integrin  $\alpha 4 \beta 1$  and CS-1 peptide specifically.

## Example 11

Construction of integrin  $\alpha 2 \cdot$  IgG heavy chain chimeric protein expression vector

[0077] The DNA fragment coding for the extracellular portion of integrin  $\alpha 2$  was divided into  $\alpha 2$ -1 and  $\alpha 2$ -2 based on the reported cDNA sequence information (Takada, Y. et al., J. Cell. Biol., 109, 397-407 (1989)) and subcloned, and they were integrated on an expression vector. At first, the RNA of human fibroblast cell line MRC-5 (ATCC CCL 171) as integrin  $\alpha 2$  expressing cell was separated, and an oligo dT cellulose column was used to purify PolyA(+)RNA. Based on it, a single stranded cDNA was synthesized and used as the template of PCR. As PCR primers, DNA oligomers of sequence Nos. 20 and 21 were synthesized for  $\alpha 2$ -1, and DNA oligomers of sequence Nos. 22 and 23, for  $\alpha 2$ -2.

5' -GCTCGAGCAAACCCAGCGCAACTACCG-3' (Sequence No. 20)

5' -ATAGTCCCCTGATGACCATTG-3' (Sequence No. 21)

5' -GATGGCTTTAATGATGTGATTG-3' (Sequence No. 22)

5' -TGTTGGTACTTCGGCTTTCTC-3' (Sequence No. 23)

[0078] The template cDNA, primers, dNTPs and Taq polymerase were mixed in a PCR buffer and PCR was performed 30 cycles by a thermal cycler (reaction conditions: 94°C 1 minute - 60°C 2 minutes - 72°C 3 minutes). The amplified DNA fragment of  $\alpha 2$ -1 was digested by restriction enzymes Xho I and EcoR I, and the DNA fragment of  $\alpha 2$ -2 was blunted at the termini by T4DNA polymerase treatment and digested by restriction enzyme EcoR I. Each fragment was purified. The two purified DNA fragments were caused to react in a phosphorylation reaction solution (50 mM Tris-HCl, 10 mM MgCl<sub>2</sub>, 25 mM DTT, 1 mM ATP, 0.1 U/ $\mu$ l T4 polynucleotide kinase (Takara), pH 8.0) at 37°C for 1 hour, and the reaction mixture was heat-treated at 68°C for 5 minutes to inactivate the enzyme. Then, IgG<sub>1</sub>SR $\alpha$  prepared in Example 1 was digested by restriction enzyme BamH I and caused to react in Klenow reaction solution (66 mM Tris-HCl, 10 mM MgCl<sub>2</sub>, 10 mM DTT, 0.2 mM dNTPs, 0.05 U/ $\mu$ l Klenow fragment (Takara), pH 7.5) at 37°C for 30 minutes, to blunt the

termini, and the reaction mixture was heat-treated at 70°C for 5 minutes to inactivate the enzyme. Furthermore, a large DNA fragment was digested by restriction enzyme Xho I, and purified. The two ( $\alpha$ 2-1 and  $\alpha$ 2-2) DNA fragments phosphorylated before were inserted into the large DNA fragment, to obtain a plasmid DNA. The obtained base sequence coding for integrin  $\alpha$ 2 • IgG heavy chain chimeric protein is shown in sequence No. 19. This plasmid (integrin  $\alpha$ 2 • IgGSR $\alpha$ ) is hereinafter called  $\alpha$ 2 • IgG heavy chain chimeric protein expression vector.

#### Example 12

Transfection of integrin  $\alpha$ 2 • IgG heavy chain chimeric protein expression vector and integrin  $\beta$ 1 • IgG heavy chain chimeric protein expression vector into animal cells, and their manifestation

[0079] The integrin  $\alpha$ 2 • IgG heavy chain chimeric protein expression vector was transfected into the integrin  $\beta$ 1 • IgG heavy chain chimeric protein producing CHO cells prepared and stabilized in Example 4, according to the lipofectin method described in Example 4. That is, integrin  $\alpha$ 2 • IgGSR $\alpha$  and pSV2neo (BRL) were mixed at 10 : 1, and the mixture was mixed with lipofectin reagent. The mixture was added dropwise to the cells. Eighteen hours after completion of dropwise addition, the mixture was cultured in a first selective medium for 2 days, and the cells were dispersed by trypsin-EDTA treatment. The cells were suspended in a second selective medium, and the suspension was disseminated into a 96-well plate. Resistant cells were selectively cultured for about 10 days. Then, the amount of integrin  $\alpha$ 2 • IgG heavy chain chimeric protein and the amount of integrin  $\beta$ 1 • IgG heavy chain chimeric protein produced in the culture supernatant were determined according to the ELISA method (described later), and a clone producing almost the same amounts of both the chimeric proteins was picked up. The clone was cloned twice according to the limiting dilution analysis, to be stabilized as a clone capable of producing  $\alpha$ 2 • IgG heavy chain- $\beta$ 1 • IgG heavy chain chimeric protein heterodimer complex.

#### Example 13

Determination of the amounts of integrin  $\alpha$ 2 • IgG heavy chain chimeric protein and integrin  $\beta$ 1 • IgG heavy chain chimeric protein by the ELISA method

[0080] Fifty microliter per well of anti-human integrin  $\alpha$ 2 antibody (Becton & Dickinson, clone P1E6) or anti-human integrin  $\beta$ 1 antibody (clone 4B4) (2  $\mu$ g/ml each) was put into a 96-well immunoplate, and allowed to stand at 4°C for 16 hours. Then, each well was washed with PBS(-) twice, blocked, and the culture supernatant of the CHO cells grown in second selective medium was properly diluted and reacted with the coated-antibody at room temperature for 1 hour. After the reaction, the surface of the plate was washed with T-PBS twice, and caused to react with biotinated anti-human IgG antibody for 1 hour and with avidin-horseradish peroxidase for 1 hour, and the reaction mixture was washed with PBS(-) twice. After completion of reaction, orthophenylenediamine was used as a substrate for color development, and the absorbance values at 490 nm were measured using a microplate reader. A clone showing a high absorbance value was selected.

#### Example 14

Purification of  $\alpha$ 2 • IgG heavy chain- $\beta$ 1 • IgG heavy chain chimeric protein heterodimer complex

(1) Culture of CHO cells and preparation of cultured supernatant solution

[0081] The CHO cells highly capable of producing  $\alpha$ 2 • IgG heavy chain- $\beta$ 1 • IgG heavy chain chimeric protein heterodimer complex were cultured in an  $\alpha$ MEM(-) medium containing 5% FBS (Ultra-low IgG grade) for 1 day, to reach semiconfluent, and they were cultured on an  $\alpha$ MEM(-) medium containing 1% FBS (Ultra-low IgG grade) for 3 days. The culture supernatant was collected, and concentrated to 1/10 volume by ultrafiltration. Then, 1M Hepes solution (pH 8.0) was added to achieve a final concentration of 5 mM, to obtain a starting solution for further purification.

(2) Protein A column chromatography

[0082] The starting solution for further purification was passed through Prosep Guard column, and applied to Prosep A column. After completion of application, it was washed with 10 times the column volume of PBS (-), and in succession, the proteins were eluted at a pH 6 to 3 gradient of 0.1M citrate buffers. The peak fraction eluted at pH 3 was collected, and 1M Tris-HCl solution (pH 8.5) was added by 0.1 volume for neutralization. The mixture was dialyzed against PBS(-).

## (3) Affinity column chromatography

[0083] According to a report (Kirchhofer, D. et al., J. Biol. Chem., 265, 615-618 (1990)), a collagen immobilized column with a collagen (Type I, Sigma) coupled to cyanogen-bromide-activated Sepharose (Sigma) was prepared. Then, the starting solution for further purification was equilibrated in a TBS buffer (150 mM NaCl, 50 mM Tris-HCl, 1 mM MgCl<sub>2</sub>, 1 mM MnCl<sub>2</sub>, pH 7.5), applied to a column, allowed to stand at room temperature for 3 hours, and washed with 10 times the column volume of a washing buffer (150 mM NaCl, 50 mM Tris-HCl, 1 mM MgCl<sub>2</sub>, 1 mM MnCl<sub>2</sub>, 100 mM octyl glucopyranoside, pH 7.5). After completion of washing, an elution buffer (20 mM EDTA, 150 mM NaCl, 50 mM Tris-HCl, 50 mM octyl glucopyranoside, pH 7.5) was used to elute the protein bound to the column. The eluate was collected and dialyzed against PBS(-).

## (4) SDS-PAGE

[0084] The eluted fraction of (3) was subjected to SDS-PAGE using 7.0% acrylamide gel under non-reducing or under reducing condition, and the gel was stained with Coomassie-blue. As a result, a band considered to be attributable to  $\alpha 2 \cdot \text{IgG heavy chain-}\beta 1 \cdot \text{IgG heavy chain chimeric protein heterodimer complex}$  was observed. Under reducing condition, two bands (185 kDa and 135 kDa) considered to be attributable to integrin  $\alpha 2 \cdot \text{IgG heavy chain chimeric protein}$  and integrin  $\beta 1 \cdot \text{IgG heavy chain chimeric protein}$  were observed. These results suggest that the eluted protein has a molecular structure considered to be  $\alpha 2 \cdot \text{IgG heavy chain-}\beta 1 \cdot \text{IgG heavy chain chimeric protein heterodimer complex}$ , and is linked by a disulfide bond between the IgG heavy chains.

## Example 15

Identification of  $\alpha 2 \cdot \text{IgG heavy chain-}\beta 1 \cdot \text{IgG heavy chain chimeric protein heterodimer complex}$ , and examination of its structural stability

[0085] The eluted protein of (3) of Example 14 was <sup>125</sup>I-labeled, and subjected to immunoprecipitation using the beads coupled with normal murine IgG, anti-human integrin  $\alpha 2$  antibody (clone P1E6) or anti-human integrin  $\beta 1$  antibody (clone 4B4) as described in Example 7, and to SDS-PAGE/autoradiography under reducing condition.

[0086] As a result, in both 1 mM MgCl<sub>2</sub> and 10 mM EDTA, from the beads of both anti-human integrin  $\alpha 2$  antibody and anti-human integrin  $\beta 1$  antibody, the same precipitation patterns expected from the structure of  $\alpha 2 \cdot \text{IgG heavy chain-}\beta 1 \cdot \text{IgG heavy chain chimeric protein heterodimer complex}$  could be obtained. These results show that the eluted protein obtained in (3) of Example 14 is certainly  $\alpha 2 \cdot \text{IgG heavy chain-}\beta 1 \cdot \text{IgG heavy chain chimeric protein heterodimer complex}$ , and with the results of (4) of Example 14 also taken into account, it is strongly suggested that the association of both the proteins is stable through a disulfide bond existing the IgG heavy chains.

## Example 16

Examination on the capability of  $\alpha 2 \cdot \text{IgG heavy chain-}\beta 1 \cdot \text{IgG heavy chain chimeric protein heterodimer complex}$  to be bound to a collagen, and its specificity

[0087] The capability of  $\alpha 2 \cdot \text{IgG heavy chain-}\beta 1 \cdot \text{IgG heavy chain chimeric protein heterodimer complex}$  to be bound to a collagen which is a ligand of integrin  $\alpha 2\beta 1$  was examined.

[0088] At first, a collagen (Cell Matrix Type 3 mg/ml) was diluted to 0.1  $\mu\text{g/ml}$  by 0.02M acetic acid solution, and put in an immunoplate by 100  $\mu\text{l/well}$ , being kept at 4°C for 16 hours. Then, the collagen solution was removed by suction, and the plate was washed with PBS(-) twice for neutralization. Heat-denaturated 1% BSA-PBS solution was put in the plate by 300  $\mu\text{l/well}$  for blocking at room temperature for 3 hours. After completion of blocking, it was rinsed With PBS (-) twice, to prepare a collagen coated plate.

[0089] The cultured supernatant of CHO (100  $\mu\text{l}$ ) containing  $\alpha 2 \cdot \text{IgG heavy chain-}\beta 1 \cdot \text{IgG heavy chain chimeric protein heterodimer complex}$  was reacted at 30°C for 3 hours. After completion of reaction, as described in Example 9, the amount of bound  $\alpha 2 \cdot \text{IgG heavy chain-}\beta 1 \cdot \text{IgG heavy chain chimeric protein heterodimer complex}$  was determined.

[0090] As a result, as shown in Fig. 4, the absorbance showing the binding of  $\alpha 2 \cdot \text{IgG heavy chain-}\beta 1 \cdot \text{IgG heavy chain chimeric protein heterodimer complex}$  to the collagen was increased. The binding was almost perfectly inhibited in the coexistence of 10  $\mu\text{g/ml}$  of anti-human integrin  $\alpha 2$  antibody (clone P1E6) and anti-human integrin  $\beta 1$  antibody (clone 4B4), or in the presence of 5 mM EDTA respectively. This result shows that  $\alpha 2 \cdot \text{IgG heavy chain-}\beta 1 \cdot \text{IgG heavy chain chimeric protein heterodimer complex}$  can be bound to a collagen like the integrin  $\alpha 2\beta 1$  existing on the surfaces of cell membranes, and furthermore that the binding is  $\alpha 2\beta 1$ -specific and that the feature of the binding that it depends on cations is retained.

## Example 17

Acquisition of a peptide capable of being bound to  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex, and evaluation of its inhibitory activity

[0091] At first,  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex purified in Example 6, or human IgG was prepared at a proper concentration by PBS (-) and was coated on a plastic plate at 4°C for 16 hours, being formed as a solid phase on a plastic plate. Then, according to a report (Cott, J. K. and Smoth, G. P., Science, 249, 386-390 (1990)), a phage peptide library in which a random six amino acid residues were cyclized by the disulfide bond of cysteine at both the ends was prepared and suspended in 0.1% BSA-containing TBS buffer. The phage peptide library was reacted with human IgG at 30°C for 3 hours, to absorb phage peptides capable of being bound to IgG. Then, the non-absorbed phases were reacted with  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex at 30°C for 3 hours, and the reaction mixture was washed with 0.1% BSA-containing TBS buffer twice to remove the phage peptides incapable of being bound to the heterodimer complex. Only the phage peptides capable of being bound were collected after elution with 0.1M glycine-hydrochloric acid (pH 2.2). After collection, the phage was amplified and the above mentioned binding operation was repeated further twice. The only the phage peptides capable of being bound to the heterodimer complex were selectively concentrated. In the final elution operation, phage peptides capable of being bound to the heterodimer complex were eluted using 10 mM EDTA and 0.1M glycine-hydrochloric acid in two steps, and the amino acid sequences of the respective peptides were analyzed. Of them, eight sequences (sequence Nos. 24 to 31) are shown in Table 1. Furthermore, they were examined using the binding assay system of Example 9, and the IC<sub>50</sub> values of the four peptide sequences showing binding inhibitory activity are shown in Table 1.

Table 1

Elution condition	Sequence								Inhibitory activity IC <sub>50</sub> (μM)	Sequence No.
EDTA	Cys*	Ile	Pro	Glu	Leu	Ile	Val	Cys*	1.2	24
	Cys*	Met	Arg	Tyr	Thr	Ser	Ala	Cys*	2.3	25
	Cys*	Glu	Trp	Met	Lys	Arg	Phe	Cys*		26
	Cys*	Tyr	Thr	Thr	Arg	Leu	Lys	Cys*		27

Glycine-hydrochloric acid	Cys*	Leu	Arg	Tyr	Ser	Val	Pro	Cys*	1.8	28
	Cys*	Ile	Val	Asn	Arg	Leu	Gly	Cys*		29
	Cys*	Gly	Leu	Gln	Ala	Leu	Pro	Cys*	10	30
	Cys*	Lys	Leu	Lys	Gly	Thr	Met	Cys*		31

Cys\* indicates a disulfide bond.

## Example 18

Acquisition of a low weight molecular compound capable of inhibiting the binding between the peptide fragment on fibronectin and  $\alpha 4 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex

[0092] Reagents and reported compounds were picked up at random, adjusted to a final concentration of 50 or 100  $\mu\text{g/ml}$ , and added to the binding determination system in Example 9. Compounds showing inhibitory activity were obtained. Of the obtained compounds, the binding inhibitory activities of the four compounds of Norethynodrel (Sigma), D-Penicillamine (Aldrich, Weigert, W. M. et al., Angew. Chem. Int. Ed. Eng., 14, 330-336 (1975)),  $\gamma$ -2-Naphthyl butyric acid (Fieser, L. F. J. Am. Chem. Soc., 70, 3197-3203 (1948)), 1-Adamantaneacetic acid (Aldrich) were shown in Table 2.

Table 2

Name of compound	Concentration ( $\mu\text{g/ml}$ )	Inhibition rate (%)
Norethynodrel	50	28
D-Penicillamine	50	51
$\gamma$ -2-Naphthyl butyric acid	100	37
1-Adamantaneacetic acid	100	65

## Example 19

Preparation of  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex liposome

[0093] A liposome was prepared according to the Martin et al.'s method (Martin, F. J. et al., Biochemistry, 20, 4229, (1981)). At first, an activated SH group was introduced into dipalmitoyl phosphatidyl ethanolamine (DPPE, Sigma) using di-crosslinking reagent N-succineimidyl 3-(2-pyridyldithio)propionate (SDPD, Sigma), to prepare pyridylthiopropionyl dipalmitoyl phosphatidyl ethanolamine (PDP-DPPE). The PDP-DPPE, dipalmitoyl phosphatidyl choline (DPPC) and cholesterol were mixed, to prepare a lipid film, and it was treated by a sonicator. Then, a filter was used to obtain a liposome uniform in diameter (PDP-DPPE liposome). Then,  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex or human IgG (Cappel) used as a negative control were dissolved in a Hepes buffer (100 mM Hepes, 150 mM NaCl, pH 8.0), and SDPD was added for reaction for 30 minutes. The reaction solution was applied to PD-10 column (Pharmacia), and eluted by 0.1M acetic acid-sodium acetate buffer (pH 5.5). To the eluate, dithiothreitol was added for treatment for 20 minutes, and the mixture was applied to PD-10 column again and eluted by a Hepes buffer (100 mM Hepes, 150 mM NaCl, pH 8.0), to obtain SDPD coupled  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex. The SDPD modified heterodimer complex and the PDP-DPPE liposome were caused to react with each other at room temperature for 24 hours, and the reaction mixture was separated by Sepharose 4B column (Sigma). From the peak fraction,  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex liposome was obtained.

[0094] The amount of  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex bound on the liposome was determined by a densitometer (ATTO) after SDS-PAGE/Coomassie staining, and adjusted to final concentration of 1 mg/ml.

## Example 20

Flow cytometry of  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex liposome

[0095]  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex liposome was dispersed in 1 mM EDTA-containing PBS(-), and caused to react with anti-human integrin  $\alpha 2$  antibody (clone P1E6) or anti-human integrin  $\beta 1$  antibody (clone 4B4) at room temperature for 30 minutes. After completion of reaction, the reaction mixture was centrifuged at 15000 rpm for 10 minutes, being followed by washing with 1 mM EDTA-containing PBS(-) and suspended into the solution again. Into the suspension, FITC labeled anti-murine IgG antibody (Cappel, 10  $\mu\text{g/ml}$ ) was added as a secondary antibody, and reacted at room temperature for 30 minutes. After completion of reaction, the reac-



tion mixture was similarly washed by centrifugation, and flow cytometry analysis (ELITE, Coulter) was performed.

[0096] As a result, the positive reactions for both the antibodies were detected, confirming that  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex was bound on the liposome.

## 5 Example 21

Binding activity of  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex to a collagen

[0097] A collagen (Cell Matrix Type, 3 mg/ml) was diluted by 0.02M acetic acid solution, and the solution was put in an immunoplate by 100  $\mu\text{l}$ /well, being kept at 4°C for 16 hours. Then, the collagen solution was removed by suction, and the plate was washed with PBS(-) twice for neutralization, and heat-denaturated 1% BSA-PBS solution was put in the plate by 300  $\mu\text{l}$ /well for blocking at room temperature for 3 hours. After completion of blocking, the plate was rinsed with PBS(-) twice, to prepare a collagen coated plate.

[0098] Normal human plasma (George King) and von Willebrand's factor deficient (severe) plasma (George Kind) were treated with anti-human IgG antibody and protein A, and dialyzed against PBS(-) for 24 hours, to remove the contained sodium citrate. In order that the Ca ion and Mg ion concentration might be a physiological concentration in the blood when used,  $\text{CaCl}_2$  and  $\text{MgCl}_2$  were added to achieve final concentrations of 1.2 mM and 0.2 mM respectively. Into the normal human plasma and von Willebrand's factor deficient plasma adjusted in cation concentration,  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex liposome or human IgG liposome was suspended to achieve protein concentrations of 1 to 100 ng/ml. Any of the suspensions was put in the collagen coated plate by 100  $\mu\text{l}$ /well. The plate was shaken by a plate shaker at 100 rpm, for reaction at room temperature for 15 minutes. After completion of reaction, the non-bound liposome was removed by washing with a PB solution (1.2 mM  $\text{CaCl}_2$ , 0.2 mM  $\text{MgCl}_2$ , 1% BSA-containing PBS, pH 7.4), and the bound liposome was immobilized by 1% glutaraldehyde-PBS at room temperature for 30 minutes. After completion of immobilization, a heat-denaturated BSA-PBS solution was used for blocking at room temperature for 1 hour. Then, as described in Example 16, it was caused to react with biotin labeled human IgG antibody used as a primary antibody and avidin labeled horseradish peroxidase used as a secondary antibody, and washed with a TBS buffer. Into it, orthophenylenediamine was added as a substrate for color development, and the absorbance at 490 nm were measured. To examine the effect of 5 mM EDTA, anti-integrin  $\alpha 2$  antibody (clone P1E6, 10  $\mu\text{g}/\text{ml}$ ) and anti-integrin  $\beta 1$  antibody(clone 4B4, 10  $\mu\text{g}/\text{ml}$ ), it was caused to react with the liposome suspension at room temperature for 15 minutes before reaction with the collagen.

[0099] The results are shown in Figs. 5 and 6. In the normal human plasma, the human IgG liposome as a negative control was not found to be bound to the collagen, but the binding of  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex liposome to the collagen was increased with the concentration dependent on manner. Also when the von Willebrand's factor deficient plasma was used, equivalent binding was detected. Furthermore, the binding to the collagen observed when 30 ng/ml of  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex liposome was added to the normal plasma was completely inhibited by adding EDTA as a cation chelating agent or the antibodies. The results show that in plasma with a physiological cation concentration,  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex liposome is bound to a collagen like platelets, and strongly suggest that it can be a substitute of adhesive platelets, and can be a reagent for monitoring the collagen exposed region. Furthermore, it is indicated that since equivalent binding activity was shown also in von Willebrand's factor deficient plasma, the liposome can also be used in the plasma with coagulation abnormality such as von Willebrand's disease.

## 45 Example 22

Analysis of collagen covering state by  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex liposome

[0100] Five microliters of a collagen solution was spotted at the center of each of the wells of a Lab-Tek chamber slide (Intermed, 8-well type, plastic) and allowed to stand for 16 hours, then washed and treated for blocking. Then, a suspension in which  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex liposome was suspended in normal human plasma to achieve a protein concentration of 30 ng/ml as described in Example 21 was put in the slide by 200  $\mu\text{l}$ /well, for reaction under the same conditions. After completion of reaction, the non-bound liposome was removed by washing with a PB buffer, and the retained was immobilized and treated for blocking. Then, it was bound to biotin labeled anti-human IgG antibody as a primary antibody and with avidin labeled horseradish peroxidase as a secondary antibody, and was washed with a TBS buffer. After completion of washing, diaminobenzidine was added for staining, to observe the covering state of the  $\alpha 2 \cdot \text{IgG}$  heavy chain- $\beta 1 \cdot \text{IgG}$  heavy chain chimeric protein heterodimer complex liposome bound on the collagen.

[0101] With the human IgG liposome, the collagen coated portion was not stained, but with  $\alpha 2 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex liposome, the collagen coated portion was entirely stained. Therefore, since the  $\alpha 2 \cdot$  IgG heavy chain- $\beta 1 \cdot$  IgG heavy chain chimeric protein heterodimer complex liposome covered the collagen coated portion only, it was strongly suggested that the liposome could be a substitute of adhesive platelets.

Industrial Availability:

[0102] The present invention provides integrin-immunoglobulin chimeric protein heterodimer complexes in which the  $\alpha$  chain and the  $\beta$  chain of an integrin are stably associated. The obtained integrin-immunoglobulin chimeric protein heterodimer complexes can be directly used as drugs, and can also be used for determining the binding between an integrin and a ligand, and searching for a substance capable of being bound to an integrin and a substance inhibiting the binding between an integrin and a ligand. They can also be used as diagnostic reagents.

[0103] Furthermore, among the heterodimer complexes, especially integrin  $\alpha 2 \beta 1$ -immunoglobulin chimeric protein heterodimer complex can be used as a substitute of platelets. Furthermore, integrin  $\alpha 2 \beta 1$ -immunoglobulin chimeric protein heterodimer complex can be used as a therapeutic or preventive agent for bleeding tendency involved in thrombocytopenia, platelet function abnormality, etc. Furthermore, it can also be used as a reagent for monitoring the exposed region of an extracellular matrix and for the targeting therapy.

## Sequence Table

5	Sequence No. 1	
	Length of sequence: 4228	
	Type of sequence: Nucleic acid	
10	Sequence	
	ATG TTC CCC ACC GAG AGC GCA TGG CTT GGG AAG CGA GGC GCG AAC CCG	48
15	Met Phe Pro Thr Glu Ser Ala Trp Leu Gly Lys Arg Gly Ala Asn Pro	
	-35 -30 -25	
	GGC CCC GAA GCT GCA CTC CGG GAG ACG GTG ATG CTG TTG CTG-TGC CTG	96
20	Gly Pro Glu Ala Ala Leu Arg Glu Thr Val Met Leu Leu Leu Cys Leu	
	-20 -15 -10	
	GGG GTC CCG ACC GGC AGG CCT TAC AAC GTG GAC ACT GAG AGC GCG CTG	144
25	Gly Val Pro Thr Gly Arg Pro Tyr Asn Val Asp Thr Glu Ser Ala Leu	
	-5 1 5	
	CTT TAC CAG GGC CCC CAC AAC ACG CTG TTC GGC TAC TCG CTC GTG CTG	192
30	Leu Tyr Gln Gly Pro His Asn Thr Leu Phe Gly Tyr Ser Val Val Leu	
	10 15 20 25	
	CAC AGC CAC GGG GCG AAC CGA TGG CTC CTA GTG GGT GCG CCC ACT GCC	240
35	His Ser His Gly Ala Asn Arg-Trp Leu Leu Val Gly Ala Pro Thr Ala	
	30 35 40	
	AAC TGG CTC GCC AAC GCT TCA GTG ATC AAT CCC GGG GCG ATT TAC AGA	288
40	Asn Trp Leu Ala Asn Ala Ser Val Ile Asn Pro Gly Ala Ile Tyr Arg	
	45 50 55	
	TGC AGG ATC GGA AAG AAT CCC GGC CAG ACG TGC GAA CAG CTC CAG CTG	336
45	Cys Arg Ile Gly Lys Asn Pro Gly Gln Thr Cys Glu Gln Leu Gln Leu	
	60 65 70	
	GGT AGC CCT AAT GGA GAA CCT TGT GGA AAG ACT TGT TTG GAA GAG AGA	384
50	Gly Ser Pro Asn Gly Glu Pro Cys Gly Lys Thr Cys Leu Glu Glu Arg	
	75 80 85	
55		

EP 0 896 002 A1

	GAC AAT CAG TCG TTC GGG GTC ACA CTT TCC AGA CAG CCA GGA GAA AAT	432
5	Asp Asn Gln Trp Leu Gly Val Thr Leu Ser Arg Gln Pro Gly Glu Asn	
	90 95 100 105	
	GGA TCC ATC GTC ACT TGT GGG CAT AGA TGG AAA AAT ATA TTT TAC ATA	480
10	Gly Ser Ile Val Thr Cys Gly His Arg Trp Lys Asn Ile Phe Tyr Ile	
	110 115 120	
	AAG AAT GAA AAT AAG CTC CCC ACT GGT GGT TGC TAT GGA GTG CCC CCT	528
15	Lys Asn Glu Asn Lys Leu Pro Thr Gly Gly Cys Tyr Gly Val Pro Pro	
	125 130 135	
	GAT TTA CGA ACA GAA CTC AGT AAA AGA ATA GCT CCG TGT TAT CAA GAT	576
20	Asp Leu Arg Thr Glu Leu Ser Lys Arg Ile Ala Pro Cys Tyr Gln Asp	
	140 145 150	
	TAT GTG AAA AAA TTT GGA GAA AAT TTT GCA TCA TGT CAA GCT GGA ATA	624
25	Tyr Val Lys Lys Phe Gly Glu Asn Phe Ala Ser Cys Gln Ala Gly Ile	
	155 160 165	
	TCC AGT TTT TAC ACA AAG GAT TTA ATT GTG ATG GGG GCC CCA GGA TCA	672
30	Ser Ser Phe Tyr Thr Lys Asp Leu Ile Val Met Gly Ala Pro Gly Ser	
	170 175 180 185	
	TCT TAC TGG ACT GGC TCT CTT TTT GTC TAC AAT ATA ACT ACA AAT AAA	720
35	Ser Tyr Trp Thr Gly Ser Leu Phe Val Tyr Asn Ile Thr Thr Asn Lys	
	190 195 200	
	TAC AAG GCT TTT TTA GAC AAA CAA AAT CAA GTA AAA TTT GGA AGT TAT	768
40	Tyr Lys Ala Phe Leu Asp Lys Gln Asn Gln Val Lys Phe Gly Ser Tyr	
	205 210 215	
	TTA GGA TAT TCA GTC GGA GCT GGT CAT TTT CGG AGC CAG CAT ACT ACC	816
45	Leu Gly Tyr Ser Val Gly Ala Gly His Phe Arg Ser Gln His Thr Thr	
	220 225 230	
	GAA GTA GTC GGA GGA GCT CCT CAA CAT GAG CAG ATT GGT AAG GCA TAT	864
50	Glu Val Val Gly Gly Ala Pro Gln His Glu Gln Ile Gly Lys Ala Tyr	
	235 240 245	

55

	ATA TTC AGC ATT GAT GAA AAA GAA CTA AAT ATC TTA CAT GAA ATG AAA	912
5	Ile Phe Ser Ile Asp Glu Lys Glu Leu Asn Ile Leu His Glu Met Lys	
	250                      255                      260                      265	
	GGT AAA AAG CTT GGA TCG TAC TTT GGA CCT TCT CTC TGT GCT GTG GAC	960
10	Gly Lys Lys Leu Gly Ser Tyr Phe Gly Ala Ser Val Cys Ala Val Asp	
	270                      275                      280	
	CTC AAT GCA GAT GGC TTC TCA GAT CTG CTC GTG GGA GCA CCC ATG CAG	1008
15	Leu Asn Ala Asp Gly Phe Ser Asp Leu Leu Val Gly Ala Pro Met Gln	
	285                      290                      295	
	AGC ACC ATC AGA GAG GAA GGA AGA GTG TTT GTG TAC ATC AAC TCT GGC	1056
20	Ser Thr Ile Arg Glu Glu Gly Arg Val Phe Val Tyr Ile Asn Ser Gly	
	300                      305                      310	
	TCG GGA GCA GTA ATG AAT GCA ATG GAA ACA AAC CTC GTT GGA AGT GAC	1104
25	Ser Gly Ala Val Met Asn Ala Met Glu Thr Asn Leu Val Gly Ser Asp	
	315                      320                      325	
	AAA TAT GCT GCA AGA TTT GGG GAA TCT ATA GTT AAT CTT GGC GAC ATT	1152
30	Lys Tyr Ala Ala Arg Phe Gly Glu Ser Ile Val Asn Leu Gly Asp Ile	
	330                      335                      340                      345	
	GAC AAT GAT GGC TTT GAA GAT GTT GCT ATC GGA GCT CCA CAA GAA GAT	1200
35	Asp Asn Asp Gly Phe Glu Asp Val Ala Ile Gly Ala Pro Gln Glu Asp	
	350                      355                      360	
	GAC TTG CAA GGT GCT ATT TAT ATT TAC AAT GGC CGT GCA GAT GGG ATC	1248
40	Asp Leu Gln Gly Ala Ile Tyr Ile Tyr Asn Gly Arg Ala Asp Gly Ile	
	365                      370                      375	
	TCG TCA ACC TTC TCA CAG AGA ATT GAA GGA CTT CAG ATC AGC AAA TCG	1296
45	Ser Ser Thr Phe Ser Gln Arg Ile Glu Gly Leu Gln Ile Ser Lys Ser	
	380                      385                      390	
	TTA AGT ATG TTT GCA CAG TCT ATA TCA GGA CAA ATT GAT GCA GAT AAT	1344
50	Leu Ser Met Phe Gly Gln Ser Ile Ser Gly Gln Ile Asp Ala Asp Asn	
	395                      400                      405	
55		

EP 0 896 002 A1

	AAT GGC TAT GTA GAT GTA GCA GTT GGT GCT TTT CGG TCT GAT TCT GCT	1392
5	Asn Gly Tyr Val Asp Val Ala Val Gly Ala Phe Arg Ser Asp Ser Ala	
	410                      415                      420                      425	
	GTC TTG CTA AGG ACA ACA CCT GTA GTA ATT GTT GAC GCT TCT TTA ACC	1440
10	Val Leu Leu Arg Thr Arg Pro Val Val Ile Val Asp Ala Ser Leu Ser	
	430                      435                      440	
	CAC CCT GAG TCA GTA AAT AGA ACG AAA TTT GAC TGT GTT GAA AAT GGA	1488
15	His Pro Glu Ser Val Asn Arg Thr Lys Phe Asp Cys Val Glu Asn Gly	
	445                      450                      455	
	TGG CCT TCT GTG TGC ATA GAT CTA ACA CTT TGT TTC TCA TAT AAG GGC	1536
20	Trp Pro Ser Val Cys Ile Asp Leu Thr Leu Cys Phe Ser Tyr Lys Gly	
	460                      465                      470	
	AAG GAA GTT CCA GGT TAC ATT GTT TTG TTT TAT AAC ATG AGT TTG GAT	1584
25	Lys Glu Val Pro Gly Tyr Ile Val Leu Phe Tyr Asn Met Ser Leu Asp	
	475                      480                      485	
	GTG AAC AGA AAG GCA GAG TCT CCA CCA AGA TTC TAT TTC TCT TCT AAT	1632
30	Val Asn Arg Lys Ala Glu Ser Pro Pro Arg Phe Tyr Phe Ser Ser Asn	
	490                      495                      500                      505	
	GGA ACT TCT GAC GTG ATT ACA GGA AGC ATA CAG GTG TCC AGC AGA GAA	1680
35	Gly Thr Ser Asp Val Ile Thr Gly Ser Ile Gln Val Ser Ser Arg Glu	
	510                      515                      520	
	GCT AAC TGT AGA ACA CAT CAA GCA TTT ATG CGG AAA GAT GTG CGG GAC	1728
40	Ala Asn Cys Arg Thr His Gln Ala Phe Met Arg Lys Asp Val Arg Asp	
	525                      530                      535	
	ATC CTC ACC CCA ATT CAG ATT GAA GCT GCT TAC CAC CTT GGT CCT CAT	1776
45	Ile Leu Thr Pro Ile Gln Ile Glu Ala Ala Tyr His Leu Gly Pro His	
	540                      545                      550	
	GTC ATC ACT AAA CGA ACT ACA GAG GAA TTC CCA CCA CTT CAG CCA ATT	1824
50	Val Ile Ser Lys Arg Ser Thr Glu Glu Phe Pro Pro Leu Gln Pro Ile	
	555                      560                      565	

55



EP 0 896 002 A1

	CTT CAG CAG AAG AAA GAA AAA CAC ATA ATG AAA AAA ACA ATA AAC TTT	1872
5	Leu Gln Gln Lys Lys Glu Lys Asp Ile Met Lys Lys Thr Ile Asn Phe	
	570                      575                      580                      585	
	GCA AGG TTT TGT CCC CAT GAA AAT TGT TCT GCT GAT TTA CAG CTT TCT	1920
10	Ala Arg Phe Cys Ala His Glu Asn Cys Ser Ala Asp Leu Gln Val Ser	
	590                      595                      600	
	GCA AAG ATT GGG TTT TTG AAG CCC CAT GAA AAT AAA ACA TAT CTT GCT	1968
15	Ala Lys Ile Gly Phe Leu Lys Pro His Glu Asn Lys Thr Tyr Leu Ala	
	605                      610                      615	
	GTT GGG AGT ATG AAG ACA TTG ATG TTG AAT GTG TCC TTG TTT AAT GCT	2016
20	Val Gly Ser Met Lys Thr Leu Met Leu Asn Val Ser Leu Phe Asn Ala	
	620                      625                      630	
	GGA GAT GAT GCA TAT GAA ACC ACT CTA CAT CTC AAA CTA CCC GTG GGT	2064
25	Gly Asp Asp Ala Tyr Glu Thr Thr Leu His Val Lys Leu Pro Val Gly	
	635                      640                      645	
	CTT TAT TTC ATT AAG ATT TTA CAG CTG GAA GAG AAG CAA ATA AAC TGT	2112
30	Leu Tyr Phe Ile Lys Ile Leu Glu Leu Glu Glu Lys Gln Ile Asn Cys	
	650                      655                      660                      665	
	GAA CTC ACA GAT AAC TCT GGC GTG GTA CAA CTT GAC TGC AGT ATT GGC	2160
35	Glu Val Thr Asp Asn Ser Gly Val Val Gln Leu Asp Cys Ser Ile Gly	
	670                      675                      680	
	TAT ATA TAT GTA GAT CAT CTC TCA AGG ATA GAT ATT AGC TTT CTC CTC	2208
40	Tyr Ile Tyr Val Asp His Leu Ser Arg Ile Asp Ile Ser Phe Leu Leu	
	685                      690                      695	
	GAT GTG AGC TCA CTC AGC AGA GCG GAA GAG GAC CTC AGT ATC ACA GTG	2256
45	Asp Val Ser Ser Leu Ser Arg Ala Glu Glu Asp Leu Ser Ile Thr Val	
	700                      705                      710	
	CAT CCT ACC TGT GAA AAT GAA GAG GAA ATG GAC AAT CTA AAG CAC ACC	2304
50	His Ala Thr Cys Glu Asn Glu Glu Glu Met Asp Asn Leu Lys His Ser	
	715                      720                      725	

55

EP 0 896 002 A1

	AGA GTG ACT GTA GCA ATA CCT TTA AAA TAT GAG GTT AAG CTG ACT GTT	2352
5	Arg Val Thr Val Ala Ile Pro Leu Lys Tyr Glu Val Lys Leu Thr Val	
	730                      735                      740                      745	
	CAT GGG TTT GTA AAC CCA ACT TCA TTT GTG TAT GGA TCA AAT GAT GAA	2400
10	His Gly Phe Val Asn Pro Thr Ser Phe Val Tyr Gly Ser Asn Asp Glu	
	750                      755                      760	
	AAT GAG CCT GAA ACG TGC ATG GTG GAG AAA ATG AAC TTA ACT TTC CAT	2448
15	Asn Glu Pro Glu Thr Cys Met Val Glu Lys Met Asn Leu Thr Phe His	
	765                      770                      775	
	GTT ATC AAC ACT GGC AAT AGT ATG GCT CCC AAT GTT AGT GTG_GAA ATA	2496
20	Val Ile Asn Thr Gly Asn Ser Met Ala Pro Asn Val Ser Val Glu Ile	
	780                      785                      790	
	ATG GTA CCA AAT TCT TTT AGC CCC CAA ACT GAT AAG CTG TTC AAC ATT	2588
25	Met Val Pro Asn Ser Phe Ser Pro Gln Thr Asp Lys Leu Phe Asn Ile	
	795                      800                      805	
	TTG GAT GTC CAG ACT ACT ACT GGA GAA TGC CAC TTT GAA AAT TAT CAA	2592
30	Leu Asp Val Gln Thr Thr Thr Gly Glu Cys His Phe Glu Asn Tyr Gln	
	810                      815                      820                      825	
	AGA GTG TGT GCA TTA GAG CAG CAA AAG AGT GCA ATG CAG ACC TTG AAA	2640
35	Arg Val Cys Ala Leu Glu Gln Gln Lys Ser Ala Met Gln Thr Leu Lys	
	830                      835                      840	
	GGC ATA GTC CGG TTC TTG TCC AAG ACT GAT AAG AGG CTA TTG TAC TGC	2688
40	Gly Ile Val Arg Phe Leu Ser Lys Thr Asp Lys Arg Leu Leu Tyr Cys	
	845                      850                      855	
	ATA AAA GCT GAT CCA CAT TGT TTA AAT TTC TTG TGT AAT TTT GGG AAA	2736
45	Ile Lys Ala Asp Pro His Cys Leu Asn Phe Leu Cys Asn Phe Gly Lys	
	860                      865                      870	
	ATG GAA AGT GGA AAA GAA GCC AGT GTT CAT ATC CAA CTG GAA GGC CGG	2784
50	Met Glu Ser Gly Lys Glu Ala Ser Val His Ile Gln Leu Glu Gly Arg	
	875                      880                      885	

55

EP 0 896 002 A1

CCA TCC ATT TTA GAA ATG GAT GAG ACT TCA CCA CTC AAG TTT GAA ATA 2832  
5 Pro Ser (Ile Leu Glu Met Asp Glu Thr Ser Ala Leu Lys Phe Glu Ile  
890 895 900 905  
AGA CCA ACA GGT TTT CCA GAG CCA AAT CCA AGA GTA ATT GAA CTA AAC 2880  
10 Arg Ala Thr Gly Phe Pro Glu Pro Asn Pro Arg Val (Ile Glu Leu Asn  
910 915 920  
AAG GAT GAG AAT GTT GCG CAT GTT CTA CTG GAA GGA CTA CAT CAT CAA 2928  
15 Lys Asp Glu Asn Val Ala His Val Leu Leu Glu Gly Leu His His Gln  
925 930 935  
AGA CCC AAA CGT TAT TTC ACG GAT CCC GAG CTGCTGGAAG CAGGCTCAGC 2978  
20 Arg Pro Lys Arg Tyr Phe Thr Asp Pro Glu  
940 945  
GCTCCTGCCT GGACGCATCC CGGCTATGCA GCGCCAGTCC AGGGCAGCAA GGCAGGCCCC 3038  
25 GTCTGCCTCT TCACCCGGAG CCTCTGCCG CCCCACCTCAT GCTCAGGGAG AGGGTCTTCT 3098  
GGCTTTTTC CAGGCTCTGG GCAGGCACAG GCTAGGTGCC CCTAACCCAG GCCCTGCACA 3158  
CAAAGGGGCA GGTGCTGGG TCAGACCTGC CAAGAGCCAT ATCCGGGAGG ACCCTGCCCC 3218  
30 TGACCTAAGC CCACCCAAA GGCCAAATC TCCACTCCCT CAGCTCGGAC ACCTTCTCTC 3278  
CTCCCAGATT CCAGTAACTC CCAATCTTCT CTCTGCA GAG CCC AAA TCT TGT GAC 3333  
Glu Pro Lys Ser Cys Asp  
35 950  
AAA ACT CAC ACA TGC CCA CCG TGC CCA GGTAAGCCAG CCCAGGCCTC 3380  
Lys Thr His Thr Cys Pro Pro Cys Pro  
40 955 960  
GCCCTCCAGC TCAAGCGGG ACAGGTGCCC TAGAGTAGCC TGCATCCAGG GACAGGCCCC 3440  
AGCCGGGTGC TGACACGTCC ACCTCCATCT CTCTCTCA GCA CCT GAA CTC CTC 3493  
45 Ala Pro Glu Leu Leu  
965  
GGG GGA CCG TCA CTC TTC CTC TTC CCC CCA AAA CCC AAG GAC ACC CTC 3541  
50 Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu  
970 975 980

55

EP 0 896 002 A1

5      ATG ATC TCC CGG ACC CCT GAG GTC ACA TGC GTG GTG GTG GAC GTG AGC      3589  
 Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser  
          985                      990                      995  
 10      CAC GAA GAC CCT GAG GTC AAG TTC AAC TGG TAC GTG GAC GGC GTG GAG      3637  
 His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu  
          1000                      1005                      1010                      1015  
 15      GTG CAT AAT GCC AAG ACA AAG CCG CGG GAG GAG CAG TAC AAC AGC ACC      3685  
 Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr  
                          1020                      1025                      1030  
 20      TAC CGG GTG GTC AGC GTC CTC ACC GTC CTG CAC CAG GAC TGG CTG AAT      3733  
 Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn  
                          1035                      1040                      1045  
 25      GGC AAG GAG TAC AAG TGC AAG GTC TCC AAC AAA GCC CTC CCA GCC CCC      3781  
 Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro  
                          1050                      1055                      1060  
 30      ATC GAG AAA ACC ATC TCC AAA GCC AAA GGTGGGACCC GTGGGGTGCG      3828  
 Ile Glu Lys Thr Ile Ser Lys Ala Lys  
                          1065                      1070  
 35      AGGGCCACAT GGACAGAGGC CGGCTCGGCC CACCCTCTGC CCTGAGAGTG ACCGCTGTAC      3888  
 CAACCTCTGT CCTACA GGG CAG CCC CGA GAA CCA CAG GTG TAC ACC CTG      3937  
                          Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
 40    1075                      1080  
 CCC CCA TCC CGG GAT GAG CTG ACC AAG AAC CAG GTC AGC CTG ACC TGC      3985  
 Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys  
 45                      1085                      1090                      1095  
 CTG GTC AAA GGC TTC TAT CCC AGC GAC ATC GCC CTG GAG TGG GAG AGC      4033  
 Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
 50                      1100                      1105                      1110                      1115  
 55

EP 0 896 002 A1

AAT GGG CAG CCG GAG AAC AAC TAC AAG ACC ACC CCT CCC CTC CTC GAT 4081  
 5 Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
 1120 1125 1130  
 TCC GAC GGC TCC TTC TTC CTC TAC AGC AAG CTC ACC GTG GAC AAG AGC 4129  
 10 Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
 1135 1140 1145  
 AGG TGG CAG CAG GGG AAC GTC TTC TCA TGC TCC GTG ATG CAT GAG GCT 4177  
 15 Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
 1150 1155 1160  
 CTG CAC AAC CAC TAC ACG CAG AAG AGC CTC TCC CTG TCT CCG-GGT AAA 4225  
 20 Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
 1165 1170 1175  
 TGA 4228  
 25  
 Sequence No. 2  
 Length of sequence: 3463  
 Type of sequence: Nucleic acid  
 30 Sequence  
 ATG AAT TTA CAA CCA ATT TTC TGG ATT GGA CTG ATC AGT TCA GTT TGC 48  
 35 Met Asn Leu Gln Pro Ile Phe Trp Ile Gly Leu Ile Ser Ser Val Cys  
 -20 -15 -10 -5  
 TGT GTG TTT GCT CAA ACA GAT GAA AAT AGA TGT TTA AAA GCA AAT GCC 96  
 40 Cys Val Phe Ala Gln Thr Asp Glu Asn Arg Cys Leu Lys Ala Asn Ala  
 1 5 10  
 AAA TCA TGT GGA GAA TGT ATA CAA GCA GGG CCA AAT TGT GGG TGG TGC 144  
 45 Lys Ser Cys Gly Glu Cys Ile Gln Ala Gly Pro Asn Cys Gly Trp Cys  
 15 20 25  
 ACA AAT TCA ACA TTT TTA CAG GAA GGA ATG CCT ACT TCT GCA CGA TGT 192  
 50 Thr Asn Ser Thr Phe Leu Gln Glu Gly Met Pro Thr Ser Ala Arg Cys  
 30 35 40  
 55

EP 0 896 002 A1

	GAT GAT TTA GAA GCC TTA AAA AAG AAG GGT TGC CCT CCA GAT GAC ATA	240
5	Asp Asp Leu Glu Ala Leu Lys Lys Lys Gly Cys Pro Pro Asp Asp Ile	
	45 50 55 60	
	GAA AAT CCC AGA GGC TCC AAA GAT ATA AAG AAA AAT AAA AAT GTA ACC	288
10	Glu Asn Pro Arg Gly Ser Lys Asp Ile Lys Lys Asn Lys Asn Val Thr	
	65 70 75	
	AAC CGT AGC AAA GGA ACA GCA GAG AAG CTC AAG CCA GAG GAT ATT CAT	336
15	Asn Arg Ser Lys Gly Thr Ala Glu Lys Leu Lys Pro Glu Asp Ile His	
	80 85 90	
	CAG ATC CAA CCA CAG CAG TTG GTT TTG CGA TTA AGA TCA GGG GAG CCA	384
20	Gln Ile Gln Pro Gln Gln Leu Val Leu Arg Leu Arg Ser Gly Glu Pro	
	95 100 105	
	CAG ACA TTT ACA TTA AAA TTC AAG AGA GCT GAA GAC TAT CCC ATT GAC	432
25	Gln Thr Phe Thr Leu Lys Phe Lys Arg Ala Glu Asp Tyr Pro Ile Asp	
	110 115 120	
	CTC TAC TAC CTT ATG GAC CTG TCT TAT TCA ATG AAA GAC GAT TTG GAG	480
30	Leu Tyr Tyr Leu Met Asp Leu Ser Tyr Ser Met Lys Asp Asp Leu Glu	
	125 130 135 140	
	AAT GTA AAA AGT CTT GGA ACA GAT CTG ATG AAT GAA ATG AGG AGG ATT	528
35	Asn Val Lys Ser Leu Gly Thr Asp Leu Met Asn Glu Met Arg Arg Ile	
	145 150 155	
	ACT TCG GAC TTC AGA ATT GGA TTT GGC TCA TTT GTG GAA AAG ACT GTG	576
40	Thr Ser Asp Phe Arg Ile Gly Phe Gly Ser Phe Val Glu Lys Thr Val	
	160 165 170	
	ATG CCT TAC ATT AGC ACA ACA CCA GCT AAG CTC AGG AAC CCT TGC ACA	624
45	Met Pro Tyr Ile Ser Thr Thr Pro Ala Lys Leu Arg Asn Pro Cys Thr	
	175 180 185	
	AGT GAA CAG AAC TGC ACC ACC CCA TTT AGC TAC AAA AAT GTG CTC AGT	672
50	Ser Glu Gln Asn Cys Thr Thr Pro Phe Ser Tyr Lys Asn Val Leu Ser	
	190 195 200	

55



EP 0 896 002 A1

	CTT ACT AAT AAA GGA GAA GTA TTT AAT GAA CTT GTT GGA AAA CAG CGC	720
5	Leu Thr Asn Lys Gly Glu Val Phe Asn Glu Leu Val Gly Lys Gln Arg	
	205                      210                      215                      220	
	ATA TCT GGA AAT TTG GAT TCT CCA GAA GGT GGT TTC GAT GCC ATC ATC	768
10	Ile Ser Gly Asn Leu Asp Ser Pro Glu Gly Gly Phe Asp Ala Ile Met	
	225                      230                      235	
	CAA GTT GCA GTT TGT GGA TCA CTG ATT GGC TGG AGG AAT GTT ACA CGG	816
15	Gln Val Ala Val Cys Gly Ser Leu Ile Gly Trp Arg Asn Val Thr Arg	
	240                      245                      250	
	CTG CTG GTG TTT TCC ACA GAT GCC GGG TTT CAC TTT GCT GGA GAT GGC	864
20	Leu Leu Val Phe Ser Thr Asp Ala Gly Phe His Phe Ala Gly Asp Gly	
	255                      260                      265	
	AAA CTT GGT GGC ATT GTT TTA CCA AAT GAT GGA CAA TGT CAC CTG GAA	912
25	Lys Leu Gly Gly Ile Val Leu Pro Asn Asp Gly Gln Cys His Leu Glu	
	270                      275                      280	
	AAT AAT ATG TAC ACA ATG AGC CAT TAT TAT GAT TAT CCT TCT ATT GCT	960
30	Asn Asn Met Tyr Thr Met Ser His Tyr Tyr Asp Tyr Pro Ser Ile Ala	
	285                      290                      295                      300	
	CAC CTT GTC CAG AAA CTG AGT GAA AAT AAT ATT CAG ACA ATT TTT GCA	1008
35	His Leu Val Gln Lys Leu Ser Glu Asn Asn Ile Gln Thr Ile Phe Ala	
	305                      310                      315	
	GTT ACT GAA GAA TTT CAG CCT GTT TAC AAG GAG CTG AAA AAC TTG ATC	1056
40	Val Thr Glu Glu Phe Gln Pro Val Tyr Lys Glu Leu Lys Asn Leu Ile	
	320                      325                      330	
	CCT AAG TCA GCA GTA GGA ACA TTA TCT GCA AAT TCT AGC AAT GTA ATT	1104
45	Pro Lys Ser Ala Val Gly Thr Leu Ser Ala Asn Ser Ser Asn Val Ile	
	335                      340                      345	
	CAG TTG ATC ATT GAT GCA TAC AAT TCC CTT TCC TCA GAA GTC ATT TTG	1152
50	Gln Leu Ile Ile Asp Ala Tyr Asn Ser Leu Ser Ser Glu Val Ile Leu	
	350                      355                      360	

55

EP 0 896 002 A1

	GAA AAC GGC AAA TTG TCA GAA GGA GTA ACA ATA AGT TAC AAA TCT TAC	1200
5	Glu Asn Gly Lys Leu Ser Glu Gly Val Thr Ile Ser Tyr Lys Ser Tyr	
	365                      370                      375                      380	
	TGC AAG AAC GGG GTG AAT GGA ACA GGG GAA AAT GGA AGA AAA TGT TCC	1248
10	Cys Lys Asn Gly Val Asn Gly Thr Gly Glu Asn Gly Arg Lys Cys Ser	
	385                      390                      395	
	AAT ATT TCC ATT GGA GAT GAG GTT CAA TTT GAA ATT AGC ATA ACT TCA	1296
15	Asn Ile Ser Ile Gly Asp Glu Val Gln Phe Glu Ile Ser Ile Thr Ser	
	400                      405                      410	
	AAT AAG TGT CCA AAA AAG GAT TCT GAC AGC TTT AAA ATT AGG CCT CTG	1344
20	Asn Lys Cys Pro Lys Lys Asp Ser Asp Ser Phe Lys Ile Arg Pro Leu	
	415                      420                      425	
	GGC TTT ACG GAG GAA GTA GAG GTT ATT CTT CAG TAC ATC TGT GAA TGT	1392
25	Gly Phe Thr Glu Glu Val Glu Val Ile Leu Gln Tyr Ile Cys Glu Cys	
	430                      435                      440	
	GAA TGC CAA AGC GAA GGC ATC CCT GAA AGT CCC AAG TGT CAT GAA GGA	1440
30	Glu Cys Gln Ser Glu Gly Ile Pro Glu Ser Pro Lys Cys His Glu Gly	
	445                      450                      455                      460	
	AAT GGG ACA TTT GAG TGT GGC GCG TGC AGG TGC AAT GAA GGG CGT GTT	1488
35	Asn Gly Thr Phe Glu Cys Gly Ala Cys Arg Cys Asn Glu Gly Arg Val	
	465                      470                      475	
	GGT AGA CAT TGT GAA TGC AGC ACA GAT GAA GTT AAC AGT GAA GAC ATC	1536
40	Gly Arg His Cys Glu Cys Ser Thr Asp Glu Val Asn Ser Glu Asp Met	
	480                      485                      490	
	GAT GCT TAC TGC AGG AAA GAA AAC AGT TCA GAA ATC TGC AGT AAC AAT	1584
45	Asp Ala Tyr Cys Arg Lys Glu Asn Ser Ser Glu Ile Cys Ser Asn Asn	
	495                      500                      505	
	GGA GAG TGC GTC TGC GGA CAG TGT GTT TGT AGC AAC AGG GAT AAT ACA	1632
50	Gly Glu Cys Val Cys Gly Gln Cys Val Cys Arg Lys Arg Asp Asn Thr	
	510                      515                      520	

55

EP 0 896 002 A1

5	AAT GAA ATT TAT TCT GCC AAA TTC TGC GAG TGT CAT AAT TTC AAC TGT Asn Glu Ile Tyr Ser Gly Lys Phe Cys Glu Cys Asp Asn Phe Asn Cys	1680
	525                      530                      535                      540	
10	GAT AGA TCC AAT GGC TTA ATT TGT GGA GGA AAT GGT GTT TGC AAG TGT Asp Arg Ser Asn Gly Leu Ile Cys Gly Gly Asn Gly Val Cys Lys Cys	1728
	545                      550                      555	
15	CGT GTC TGT GAG TGC AAC CCC AAC TAC ACT GGC AGT GCA TGT GAC TGT Arg Val Cys Glu Cys Asn Pro Asn Tyr Thr Gly Ser Ala Cys Asp Cys	1776
	560                      565                      570	
20	TCT TTG GAT ACT AGT ACT TGT GAA GCC AGC AAC GGA CAG ATC TGC AAT Ser Leu Asp Thr Ser Thr Cys Glu Ala Ser Asn Gly Gln Ile Cys Asn	1824
	575                      580                      585	
25	GGC CGG GGC ATC TGC GAG TGT GGT GTC TGT AAG TGT ACA GAT CCG AAG Gly Arg Gly Ile Cys Glu Cys Gly Val Cys Lys Cys Thr Asp Pro Lys	1872
	590                      595                      600	
30	TTT CAA GGG CAA ACG TGT GAG ATG TGT CAG ACC TGC CTT GGT GTC TGT Phe Gln Gly Gln Thr Cys Glu Met Cys Gln Thr Cys Leu Gly Val Cys	1920
	605                      610                      615                      620	
35	GCT GAG CAT AAA GAA TGT GTT CAG TGC AGA GCC TTC AAT AAA GGA GAA Ala Glu His Lys Glu Cys Val Gln Cys Arg Ala Phe Asn Lys Gly Glu	1968
	625                      630                      635	
40	AAG AAA GAC ACA TGC ACA CAG GAA TGT TCC TAT TTT AAC ATT ACC AAG Lys Lys Asp Thr Cys Thr Gln Glu Cys Ser Tyr Phe Asn Ile Thr Lys	2016
	640                      645                      650	
45	GTA GAA AGT CCG GAC AAA TTA CCC CAG CCG GTC CAA CCT GAT CCT GTC Val Glu Ser Arg Asp Lys Leu Pro Gln Pro Val Gln Pro Asp Pro Val	2064
	655                      660                      665	
50	TCC CAT TGT AAG GAG AAG GAT GTT GAC GAC TGT TGC TTC TAT TTT ACG Ser His Cys Lys Glu Lys Asp Val Asp Asp Cys Trp Phe Tyr Phe Thr	2112
	670                      675                      680	

55

EP 0 896 002 A1

5 TAT TCA GTG AAT GGG AAC AAC GAG GTC ATG GTT CAT GTT GTG GAG AAT 2160  
 Tyr Ser Val Asn Gly Asn Asn Glu Val Met Val His Val Val Glu Asn  
 685 690 695 700  
 10 CCA GAG TGT CCC ACT GGT CCA GAG GAT CCC GAG CTGCTGGAAG CAGGCTCAGC 2213  
 Pro Glu Cys Pro Thr Gly Pro Glu Asp Pro Glu  
 705 710  
 15 GCTCCTGCCT GGACGCATCC CGGCTATGCA GCGCCAGTCC AGGGCAGCAA GGCAGGCCCC 2273  
 GTCTGCCTCT TCACCCGGAG CCTCTGCCCG CCCCACTCAT GCTCAGGGAG AGGGTCTTCT 2333  
 GGCTTTTTC CAGGCTCTGG GCAGGCACAG GCTAGGTGCC CCTAACCCAG GCCCTGCACA 2393  
 CAAAGGGGCA GGTGCTGGGC TCAGACCTGC CAAGAGCCAT ATCCGGGAGG ACCCTGCCCC 2453  
 20 TGACCTAAGC CCACCCAAA GGCCAACTC TCCACTCCCT CAGCTCGGAC ACCTTCTCTC 2513  
 CTCCCAGATT CCAGTAACTC CCAATCTTCT CTCTGCA GAG CCC AAA TCT TGT GAC 2568  
 Glu Pro Lys Ser Cys Asp  
 25 715  
 AAA ACT CAC ACA TGC CCA CCG TGC CCA GGTAAGCCAG CCCAGGCCTC 2615  
 Lys Thr His Thr Cys Pro Pro Cys Pro  
 30 720 725  
 GCCCTCCAGC TCAAGGCGGG ACAGGTGCCC TAGAGTAGCC TGCATCCAGG GACAGGCCCC 2675  
 AGCCGGGTGC TGACACGTCC ACCTCCATCT CTTCCTCA GCA CCT GAA CTC CTG 2728  
 35 Ala Pro Glu Leu Leu  
 730  
 40 GGG GGA CCG TCA GTC TTC CTC TTC CCC CCA AAA CCC AAG GAC ACC CTC 2776  
 Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu  
 735 740 745  
 45 ATG ATC TCC CGG ACC CCT GAG GTC ACA TGC GTG GTG GTG GAC GTG ACC 2824  
 Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser  
 750 755 760  
 50 CAC GAA GAC CCT GAG GTC AAG TTC AAC TGG TAC GTG GAC GGC GTG GAG 2872  
 His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu  
 765 770 775  
 55

EP 0 896 002 A1

5 GTG CAT AAT GCC AAG ACA AAG CCG CGG GAG GAG CAG TAC AAC AGC ACC 2920  
 Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr  
 780 785 790 795  
 10 TAC CGG GTG GTC AGC GTC CTC ACC GTC CTC CAC CAG GAC TGG CTG AAT 2968  
 Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn  
 800 805 810  
 15 GGC AAG GAG TAC AAG TGC AAG GTC TCC AAC AAA GCC CTC CCA GCC CCC 3016  
 Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro  
 815 820 825  
 20 ATC GAG AAA ACC ATC TCC AAA GCC AAA GGTGGGACCC GTGGGGTGG 3063  
 Ile Glu Lys Thr Ile Ser Lys Ala Lys  
 830 835  
 25 AGGGCCACAT GGACAGAGGC CGGCTCGGCC CACCCTCTGC CCTGAGAGTG ACCGCTGTAC 3123  
 CAACCTCTGT CCTACA GGG CAG CCC CGA GAA CCA CAG GTG TAC ACC CTG 3172  
 Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu  
 30 840 845  
 CCC CCA TCC CGG GAT GAG CTG ACC AAG AAC CAG GTC AGC CTG ACC TGC 3220  
 Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys  
 35 850 855 860  
 CTG GTC AAA GGC TTC TAT CCC AGC GAC ATC GCC GTG GAG TGG GAG AGC 3268  
 Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser  
 40 865 870 875  
 AAT GGG CAG CCG GAG AAC AAC TAC AAG ACC ACG CCT CCC GTG CTG GAT 3316  
 Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp  
 45 880 885 890 895  
 TCC GAC GGC TCC TTC TTC CTC TAC AGC AAG CTC ACC GTG GAC AAC AGC 3364  
 Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser  
 50 900 905 910

55

EP 0 896 002 A1

AGG TGG CAG CAG GCG AAC GTC TTC TCA TGC TCC GTG ATG CAT GAG GCT 3412  
 5 Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala  
 915 920 925  
 CTG CAC AAC CAC TAC ACG CAG AAG AGC CTC TCC CTG TCT CCG GGT AAA 3460  
 10 Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
 930 935 940  
 TGA 3463

15

Sequence No. 3

Length of sequence: 13

20

Type of sequence: Amino acid

Topology: Linear

Kind of sequence: Peptide

25

Sequence

Cys Leu His Gly Pro Glu Ile Leu Asp Val Pro Ser Thr

1 5 10

30

Sequence No. 4

Length of sequence: 31

35

Type of sequence: Nucleic acid

Number of strands: Single

Topology: Linear

40

Kind of sequence: Other nucleic acid, synthetic DNA

Sequence

GCGGATCCCG AGCTGCTGGA AGCAGGCTCA G

31

45

Sequence No. 5

Length of sequence: 27

50

Type of sequence: Nucleic acid

Number of strands: Single

55

Topology: Linear

5 Kind of sequence: Other nucleic acid, synthetic DNA  
Sequence

CCTCTAGACG GCCGTCGCAC TCATTTA

27

10

Sequence No. 6

Length of sequence: 73

15 Type of sequence: Nucleic acid

Number of strands: Single

Topology: Linear

20 Kind of sequence: Other nucleic acid, synthetic DNA  
Sequence

CTAGACCACC ATGTTCCCA CCGAGAGCGC ATGGCTTGGG AAGCGAGGCG CGAACCCGGG  
25 CCCCAGAGCT GCA

73

Sequence No. 7

30 Length of sequence: 65

Type of sequence: Nucleic acid

Number of strands: Single

35 Topology: Linear

Kind of sequence: Other nucleic acid, synthetic DNA  
Sequence

40 GCTTCGGGGC CCGGTTCCG GCCTCGCTTC CCAAGCCATG CGCTCTCGGT GGGGAACATG  
GTGCT

65

45

Sequence No. 8

Length of sequence: 51

Type of sequence: Nucleic acid

50 Number of strands: Single

Topology: Linear

55

Kind of sequence: Other nucleic acid, synthetic DNA  
Sequence

5 CTCCGGGAGA CGGTGATGCT GTTGCTGTGC CTGGGGGTCC CGACCGGCAG G 51

Sequence No. 9

10 Length of sequence: 55

Type of sequence: Nucleic acid

Number of strands: Single

15 Topology: Straight chain

Kind of sequence: Other nucleic acid, synthetic DNA  
Sequence

20 CCTGCCGGTC GGGACCCCA GGCACAGCAA CAGCATCACC GTCTCCCGGA GTCGA 55

Sequence No. 10

25 Length of sequence: 37

Type of sequence: Nucleic acid

Number of strands: Single

30 Topology: Linear

Kind of sequence: Other nucleic acid, synthetic DNA  
Sequence

35 CACTGCAGGC AGGCCTTACA ACGTGGACAC TGAGAGC 37

Sequence No. 11

40 Length of sequence: 22

Type of sequence: Nucleic acid

Number of strands: Single

45 Topology: Linear

Kind of sequence: Other nucleic acid, synthetic DNA  
Sequence

50 GCAGAAACCT GTAAATCAGC AG 22

55



Sequence No. 12

Length of sequence: 22

Type of sequence: Nucleic acid

Number of strands: Single

Topology: Linear

Kind of sequence: Other nucleic acid, synthetic DNA  
Sequence

GCATTATGC GGAAAGATGT GC

22

Sequence No. 13

Length of sequence: 29

Type of sequence: Nucleic acid

Number of strands: Single

Topology: Linear

Kind of sequence: Other nucleic acid, synthetic DNA  
Sequence

CGGGATCCGT GAAATAACGT TTGGGTCTT

29

Sequence No. 14

Length of sequence: 22

Type of sequence: Nucleic acid

Number of strands: Single

Topology: Linear

Kind of sequence: Other nucleic acid, synthetic DNA  
Sequence

GCGGAAAAGA TGAATTACA AC

22

Sequence No. 15

Length of sequence: 27

Type of sequence: Nucleic acid  
 Number of strands: Single  
 Topology: Linear  
 Kind of sequence: Other nucleic acid, synthetic DNA  
 Sequence

GTGGGATCCT CTGGACCACT GGCACAC

27

Sequence No. 16  
 Length of sequence: 10  
 Type of sequence: Amino acid  
 Topology: Linear  
 Kind of sequence: Peptide  
 Sequence

Gly Pro Glu Ile Leu Asp Val Pro Ser Thr  
 1 5 10

Sequence No. 17  
 Length of sequence: 10  
 Type of sequence: Amino acid  
 Topology: Linear  
 Kind of sequence: Peptide  
 Sequence

Gly Pro Glu Ile Leu Glu Val Pro Ser Thr  
 1 5 10

Sequence No. 18  
 Length of sequence: 6  
 Type of sequence: Amino acid  
 Topology: Linear  
 Kind of sequence: Peptide

## Sequence

5 Gly Arg Gly Asp Ser Pro

1 5

Sequence No. 19

10 Length of sequence: 4675

Type of sequence: Nucleic acid

## Sequence

15 ATG GGG CCA GAA CGG ACA GGG GCC GCG CCG CTG CCG CTG CTG CTG GTG 48  
Met Gly Pro Glu Arg Thr Gly Ala Ala Pro Leu Pro Leu Leu Leu Val  
-25 -20 -15  
20 TTA GCG CTC AGT CAA GGC ATT TTA AAT TGT TGT TTG GCC TAC AAT GTT 96  
Leu Ala Leu Ser Gln Gly Ile Leu Asn Cys Cys Leu Ala Tyr Asn Val  
-10 -5 1  
25 GGT CTC CCA GAA GCA AAA ATA TTT TCC GGT CCT TCA AGT GAA CAG TTT 114  
Gly Leu Pro Glu Ala Lys Ile Phe Ser Gly Pro Ser Ser Glu Gln Phe  
5 10 15  
30 GGG TAT GCA GTG CAG CAG TTT ATA AAT CCA AAA GGC AAC TGG TTA CTG 192  
Gly Tyr Ala Val Gln Gln Phe Ile Asn Pro Lys Gly Asn Trp Leu Leu  
35 20 25 30 35  
GTT GGT TCA CCC TGG AGT GGC TTT CCT GAG AAC CGA ATG GGA GAT GTG 240  
Val Gly Ser Pro Trp Ser Gly Phe Pro Glu Asn Arg Met Gly Asp Val  
40 40 45 50  
TAT AAA TGT CCT GTT GAC CTA TCC ACT GCC ACA TGT GAA AAA CTA AAT 288  
Tyr Lys Cys Pro Val Asp Leu Ser Thr Ala Thr Cys Glu Lys Leu Asn  
45 55 60 65  
TTG CAA ACT TCA ACA AGC ATT CCA AAT GTT ACT GAG ATG AAA ACC AAC 336  
Leu Gln Thr Ser Thr Ser Ile Pro Asn Val Thr Glu Met Lys Thr Asn  
50 70 75 80

55

EP 0 896 002 A1

	ATG AGC CTC GGC TTG ATC CTC ACC AGG AAC ATG GGA ACT GGA GGT TTT	384
5	Met Ser Leu Gly Leu Ile Leu Thr Arg Asn Met Gly Thr Gly Gly Phe	
	85 90 95	
	CTC ACA TGT GGT CCT CTG TGG GCA CAG CAA TGT GGC AAT CAG TAT TAC	432
10	Leu Thr Cys Gly Pro Leu Trp Ala Gln Gln Cys Gly Asn Gln Tyr Tyr	
	100 105 110 115	
	ACA ACG GGT GTG TGT TCT GAC ATC AGT CCT GAT TTT CAG CTC TCA GCC	480
15	Thr Thr Gly Val Cys Ser Asp Ile Ser Pro Asp Phe Gln Leu Ser Ala	
	120 125 130	
	AGC TTC TCA CCT GCA ACT CAG CCC TGC CCT TCC CTC ATA GAT GTT GTG	528
20	Ser Phe Ser Pro Ala Thr Gln Pro Cys Pro Ser Leu Ile Asp Val Val	
	135 140 145	
	GTT GTG TGT GAT GAA TCA AAT AGT ATT TAT CCT TGG GAT GCA GTA AAG	576
25	Val Val Cys Asp Glu Ser Asn Ser Ile Tyr Pro Trp Asp Ala Val Lys	
	150 155 160	
	AAT TTT TTG GAA AAA TTT GTA CAA GGC CTT GAT ATA GGC CCC ACA AAG	624
30	Asn Phe Leu Glu Lys Phe Val Gln Gly Leu Asp Ile Gly Pro Thr Lys	
	165 170 175	
	ACA CAG GTG GGG TTA ATT CAG TAT GCC AAT AAT CCA AGA GTT GTG TTT	672
35	Thr Gln Val Gly Leu Ile Gln Tyr Ala Asn Asn Pro Arg Val Val Phe	
	180 185 190 195	
	AAC TTG AAC ACA TAT AAA ACC AAA GAA GAA ATG ATT GTA GCA ACA TCC	720
40	Asn Leu Asn Thr Tyr Lys Thr Lys Glu Glu Met Ile Val Ala Thr Ser	
	200 205 210	
	CAG ACA TCC CAA TAT GGT GGG GAC CTC ACA AAC ACA TTC GGA GCA ATT	768
45	Gln Thr Ser Gln Tyr Gly Gly Asp Leu Thr Asn Thr Phe Gly Ala Ile	
	215 220 225	
	CAA TAT GCA AGA AAA TAT GCC TAT TCA GCA GCT TCT GGT GGC CGA CGA	816
50	Gln Tyr Ala Arg Lys Tyr Ala Tyr Ser Ala Ala Ser Gly Gly Arg Arg	
	230 235 240	
55		

EP 0 896 002 A1

	AGT GCT ACG AAA GTA ATG CTA GTT CTA ACT GAC GGT GAA TCA CAT GAT	864
5	Ser Ala Thr Lys Val Met Val Val Val Thr Asp Gly Glu Ser His Asp	
	245 250 255	
	GGT TCA ATG TTG AAA GCT GTG ATT GAT CAA TGC AAC CAT GAC AAT ATA	912
10	Gly Ser Met Leu Lys Ala Val Ile Asp Gln Cys Asn His Asp Asn Ile	
	260 265 270 275	
	CTG AGG TTT GGC ATA GCA GTT CTT GGG TAC TTA AAC AGA AAC GCC CTT	960
15	Leu Arg Phe Gly Ile Ala Val Leu Gly Tyr Leu Asn Arg Asn Ala Leu	
	280 285 290	
	GAT ACT AAA AAT TTA ATA AAA GAA ATA AAA GCG ATC GCT AGT ATT CCA	1008
20	Asp Thr Lys Asn Leu Ile Lys Glu Ile Lys Ala Ile Ala Ser Ile Pro	
	295 300 305	
	ACA GAA AGA TAC TTT TTC AAT GTG TCT GAT GAA GCA GCT CTA CTA GAA	1056
25	Thr Glu Arg Tyr Phe Phe Asn Val Ser Asp Glu Ala Ala Leu Leu Glu	
	310 315 320	
	AAG GCT GGC ACA TTA GGA GAA CAA ATT TTC AGC ATT GAA GGT ACT GTT	1104
30	Lys Ala Gly Thr Leu Gly Glu Gln Ile Phe Ser Ile Glu Gly Thr Val	
	325 330 335	
	CAA GGA GGA GAC AAC TTT CAG ATG GAA ATG TCA CAA GTG GGA TTC AGT	1152
35	Gln Gly Gly Asp Asn Phe Gln Met Glu Met Ser Gln Val Gly Phe Ser	
	340 345 350 355	
	GCA GAT TAC TCT TCT CAA AAT GAT ATT CTG ATG CTG GGT GCA GTG GGA	1200
40	Ala Asp Tyr Ser Ser Gln Asn Asp Ile Leu Met Leu Gly Ala Val Gly	
	360 365 370	
	GCT TTT GGC TGG AGT GGC ACC ATT GTC CAG AAG ACA TCT CAT GGC CAT	1248
45	Ala Phe Gly Trp Ser Gly Thr Ile Val Gln Lys Thr Ser His Gly His	
	375 380 385	
	TTG ATC TTT CCT AAA CAA GCC TTT GAC CAA ATT CTG CAG GAC AGA AAT	1296
50	Leu Ile Phe Pro Lys Gln Ala Phe Asp Gln Ile Leu Gln Asp Arg Asn	
	390 395 400	
55		

EP 0 896 002 A1

	CAC ACT TCA TAT TTA GGT TAC TCT GTG GCT GCA ATT TCT ACT GGA GAA	1344
5	His Ser Ser Tyr Leu Gly Tyr Ser Val Ala Ala Ile Ser Thr Gly Glu	
	405 410 415	
	AGC ACT CAC TTT GTT GCT GCT CCT CGG GCA AAT TAT ACC GGC CAG	1392
10	Ser Thr His Phe Val Ala Gly Ala Pro Arg Ala Asn Tyr Thr Gly Gln	
	420 425 430 435	
	ATA GTG CTA TAT AGT GTG AAT GAG AAT GGC AAT ATC ACG GTT ATT CAG	1440
15	Ile Val Leu Tyr Ser Val Asn Glu Asn Gly Asn Ile Thr Val Ile Gln	
	440 445 450	
	GCT CAC CGA GGT GAC CAG ATT GGC TCC TAT TTT GGT AGT GTG CTG TGT	1488
20	Ala His Arg Gly Asp Gln Ile Gly Ser Tyr Phe Gly Ser Val Leu Cys	
	455 460 465	
	TCA GTT GAT GTG GAT AAA GAC ACC ATT ACA GAC GTG CTC TTG GTA GGT	1536
25	Ser Val Asp Val Asp Lys Asp Thr Ile Thr Asp Val Leu Leu Val Gly	
	470 475 480	
	GCA CCA ATG TAC ATG AGT GAC CTA AAG AAA GAG GAA GGA AGA GTC TAC	1584
30	Ala Pro Met Tyr Met Ser Asp Leu Lys Lys Glu Glu Gly Arg Val Tyr	
	485 490 495	
	CTG TTT ACT ATC AAA AAG GGC ATT TTG GGT CAG CAC CAA TTT CTT GAA	1632
35	Leu Phe Thr Ile Lys Lys Gly Ile Leu Gly Gln His Gln Phe Leu Glu	
	500 505 510 515	
	GGC CCC GAG GGC ATT GAA AAC ACT CGA TTT GGT TCA GCA ATT GCA GCT	1680
40	Gly Pro Glu Gly Ile Glu Asn Thr Arg Phe Gly Ser Ala Ile Ala Ala	
	520 525 530	
	CTT TCA GAC ATC AAC ATG GAT GGC TTT AAT GAT GTG ATT GTT GGT TCA	1728
45	Leu Ser Asp Ile Asn Met Asp Gly Phe Asn Asp Val Ile Val Gly Ser	
	535 540 545	
	CCA CTA GAA AAT CAG AAT TCT GGA GCT GTA TAC ATT TAC AAT GGT CAT	1776
50	Pro Leu Glu Asn Gln Asn Ser Gly Ala Val Tyr Ile Tyr Asn Gly His	
	550 555 560	

55

EP 0 896 002 A1

	CAG GGC ACT ATC CGC ACA AAG TAT TCC CAG AAA ATC TTG GGA TCC GAT	1824
5	Gln Gly Thr Ile Arg Thr Lys Tyr Ser Gln Lys Ile Leu Gly Ser Asp	
	565                      570                      575	
	GGA GCC TTT AGG AGC CAT CTC CAG TAC TTT GGG AGG TCC TTG GAT GGC	1872
10	Gly Ala Phe Arg Ser His Leu Gln Tyr Phe Gly Arg Ser Leu Asp Gly	
	580                      585                      590                      595	
	TAT GGA GAT TTA AAT GGG GAT TCC ATC ACC GAT GTG TCT ATT GGT GCC	1920
15	Tyr Gly Asp Leu Asn Gly Asp Ser Ile Thr Asp Val Ser Ile Gly Ala	
	600                      605                      610	
	TTT GGA CAA GTG GTT CAA CTC TGG TCA CAA AGT ATT GCT GAT GTA GCT	1968
20	Phe Gly Gln Val Val Gln Leu Trp Ser Gln Ser Ile Ala Asp Val Ala	
	615                      620                      625	
	ATA GAA GCT TCA TTC ACA CCA GAA AAA ATC ACT TTG GTC AAC AAG AAT	2016
25	Ile Glu Ala Ser Phe Thr Pro Glu Lys Ile Thr Leu Val Asn Lys Asn	
	630                      635                      640	
	GCT CAG ATA ATT CTC AAA CTC TGC TTC AGT GCA AAG TTC AGA CCT ACT	2064
30	Ala Gln Ile Ile Leu Lys Leu Cys Phe Ser Ala Lys Phe Arg Pro Thr	
	645                      650                      655	
	AAG CAA AAC AAT CAA GTG GCC ATT GTA TAT AAC ATC ACA CTT GAT GCA	2112
35	Lys Gln Asn Asn Gln Val Ala Ile Val Tyr Asn Ile Thr Leu Asp Ala	
	660                      665                      670                      675	
	GAT GGA TTT TCA TCC AGA GTA ACC TCC AGG GGG TTA TTT AAA GAA AAC	2160
40	Asp Gly Phe Ser Ser Arg Val Thr Ser Arg Gly Leu Phe Lys Glu Asn	
	680                      685                      690	
	AAT GAA AGG TGC CTG CAG AAG AAT ATG GTA GTA AAT CAA GCA CAG AGT	2208
45	Asn Glu Arg Cys Leu Gln Lys Asn Met Val Val Asp Gln Ala Gln Ser	
	695                      700                      705	
	TGC CCC GAG CAC ATC ATT TAT ATA CAG GAG CCC TCT GAT GTT GTC AAC	2256
50	Cys Pro Glu His Ile Ile Tyr Ile Gln Glu Pro Ser Asp Val Val Asn	
	710                      715                      720	

55

EP 0 896 002 A1

	TCT TTG GAT TTG CGT GTG GAC ATC AGT CTG GAA AAC CCT GCC ACT AGC	2304
5	Ser Leu Asp Leu Arg Val Asp Ile Ser Leu Glu Asn Pro Gly Thr Ser	
	725 730 735	
	CCT GCC CTT GAA GCC TAT TCT GAG ACT GCC AAG GTC TTC AGT ATT CCT	2352
10	Pro Ala Leu Glu Ala Tyr Ser Glu Thr Ala Lys Val Phe Ser Ile Pro	
	740 745 750 755	
	TTC CAC AAA GAC TGT GGT GAG GAT GGA CTT TGC ATT TCT GAT CTA GTC	2400
15	Phe His Lys Asp Cys Gly Glu Asp Gly Leu Cys Ile Ser Asp Leu Val	
	760 765 770	
	CTA GAT GTC CGA CAA ATA CCA GCT GCT CAA GAA CAA CCC TTT ATT GTC	2448
20	Leu Asp Val Arg Gln Ile Pro Ala Ala Gln Glu Gln Pro Phe Ile Val	
	775 780 785	
	AGC AAC CAA AAC AAA AGG TTA ACA TTT TCA GTA ACA CTG AAA AAT AAA	2496
25	Ser Asn Gln Asn Lys Arg Leu Thr Phe Ser Val Thr Leu Lys Asn Lys	
	790 795 800	
	AGG GAA AGT GCA TAC AAC ACT GGA ATT GTT GTT GAT TTT TCA GAA AAC	2544
30	Arg Glu Ser Ala Tyr Asn Thr Gly Ile Val Val Asp Phe Ser Glu Asn	
	805 810 815	
	TTG TTT TTT GCA TCA TTC TCC CTA CCG GTT GAT GGG ACA GAA GTA ACA	2592
35	Leu Phe Phe Ala Ser Phe Ser Leu Pro Val Asp Gly Thr Glu Val Thr	
	820 825 830 835	
	TGC CAG GTC GCT GCA TCT CAG AAG TCT GTT GCC TGC GAT GTA GGC TAC	2640
40	Cys Gln Val Ala Ala Ser Gln Lys Ser Val Ala Cys Asp Val Gly Tyr	
	840 845 850	
	CCT GCT TTA AAG AGA GAA CAA CAG GTG ACT TTT ACT ATT AAC TTT GAC	2688
45	Pro Ala Leu Lys Arg Glu Gln Gln Val Thr Phe Thr Ile Asn Phe Asp	
	855 860 865	
	TTC AAT CTT CAA AAC CTT CAG AAT CAG GCG TCT CTC AGT TTC CAA GCC	2736
50	Phe Asn Leu Gln Asn Leu Gln Asn Gln Ala Ser Leu Ser Phe Gln Ala	
	870 875 880	

55



EP 0 896 002 A1

5 TTA AGT GAA AGC CAA GAA GAA AAC AAG GCT GAT AAT TTG GTC AAC CTC 2784  
Leu Ser Glu Ser Gln Glu Glu Asn Lys Ala Asp Asn Leu Val Asn Leu  
885 890 895

10 AAA ATT CCT CTC CTG TAT GAT GCT GAA ATT CAC TTA ACA AGA TCT ACC 2832  
Lys Ile Pro Leu Leu Tyr Asp Ala Glu Ile His Leu Thr Arg Ser Thr  
900 905 910 915

15 AAC ATA AAT TTT TAT GAA ATC TCT TCG GAT GGG AAT GTT CCT TCA ATC 2880  
Asn Ile Asn Phe Tyr Glu Ile Ser Ser Asp Gly Asn Val Pro Ser Ile  
920 925 930

20 GTG CAC AGT TTT GAA GAT GTT GGT CCA AAA TTC ATC TTC TCC CTG AAG 2928  
Val His Ser Phe Glu Asp Val Gly Pro Lys Phe Ile Phe Ser Leu Lys  
935 940 945

25 GTA ACA ACA GGA AGT GTT CCA GTA AGC ATG GCA ACT GTA ATC ATC CAC 2976  
Val Thr Thr Gly Ser Val Pro Val Ser Met Ala Thr Val Ile Ile His  
950 955 960

30 ATC CCT CAG TAT ACC AAA GAA AAG AAC CCA CTG ATG TAC CTA ACT GGG 3024  
Ile Pro Gln Tyr Thr Lys Glu Lys Asn Pro Leu Met Tyr Leu Thr Gly  
965 970 975

35 GTG CAA ACA GAC AAG GCT GGT GAC ATC AGT TGT AAT GCA GAT ATC AAT 3072  
Val Gln Thr Asp Lys Ala Gly Asp Ile Ser Cys Asn Ala Asp Ile Asn  
980 985 990 995

40 CCA CTG AAA ATA GGA CAA ACA TCT TCT TCT GTA TCT TTC AAA AGT GAA 3120  
Pro Leu Lys Ile Gly Gln Thr Ser Ser Ser Val Ser Phe Lys Ser Glu  
1000 1005 1010

45 AAT TTC AGG CAC ACC AAA GAA TTG AAC TGC AGA ACT GCT TCC TGT AGT 3168  
Asn Phe Arg His Thr Lys Glu Leu Asn Cys Arg Thr Ala Ser Cys Ser  
1015 1020 1025

50 AAT GTT ACC TGC TGG TTG AAA GAC GTT CAC ATG AAA GGA GAA TAC TTT 3216  
Asn Val Thr Cys Trp Leu Lys Asp Val His Met Lys Gly Glu Tyr Phe  
1030 1035 1040

55

EP 0 896 002 A1

5 GTT AAT GTG ACT ACC AGA ATT TGG AAC GGG ACT TTC GCA TCA TCA ACC 3264  
 Val Asn Val Thr Thr Arg Ile Trp Asn Gly Thr Phe Ala Ser Ser Thr  
 1045 1050 1055  
 10 TTC CAG ACA GTA CAG CTA ACC GCA GCT GCA GAA ATC AAC ACC TAT AAC 3312  
 Phe Gln Thr Val Gln Leu Thr Ala Ala Ala Glu Ile Asn Thr Tyr Asn  
 1060 1065 1070 1075  
 15 CCT CAG ATA TAT GTG ATT GAA GAT AAC ACT GTT ACG ATT CCC CTG ATG 3360  
 Pro Glu Ile Tyr Val Ile Glu Asp Asn Thr Val Thr Ile Pro Leu Met  
 1080 1085 1090  
 20 ATA ATG AAA CCT GAT GAG AAA GCC GAA GTA CCA ACA GAT CCC GAG 3405  
 Ile Met Lys Pro Asp Glu Lys Ala Glu Val Pro Thr Asp Pro Glu  
 1095 1100 1105  
 25 CTGCTGGAAG CAGGCTCAGC GCTCCTGCCT GGACGCATCC CGGCTATGCA GCGCCAGTCC 3465  
 AGGGCAGCAA GGCAGGCCCC GTCTGCCTCT TCACCCGGAG CCTCTGCCCG CCCCACTCAT 3525  
 GCTCAGGGAG AGGGTCTTCT GGCTTTTTC CAGGCTCTGG GCAGGCACAG GCTAGGTGCC 3585  
 CCTAACCCAG GCCCTGCACA CAAAGGGGCA GGTGCTGGGC TCAGACCTGC CAAGAGCCAT 3645  
 30 ATCCGGGAGG ACCCTGCCCC TGACCTAAGC CCACCCCAAA GGCCAAACTC TCCACTCCCT 3705  
 CAGCTCGGAC ACCTTCTCTC CTCCCAGATT CCAGTAACTC CCAATCTTCT CTCTGCA 3762  
 GAG CCC AAA TCT TGT GAC AAA ACT CAC ACA TGC CCA CCG TGC CCA 3807  
 35 Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro  
 1110 1115 1120  
 40 GGTAAGCCAG CCCAGGCCCTC GCCCTCCAGC TCAAGGCGGG ACAGGTGCCC TAGAGTAGCC 3867  
 TGCATCCAGG GACAGGCCCC AGCCGGGTGC TGACACGTCC ACCTCCATCT CTTCCTCA 3925  
 GCA CCT GAA CTC CTG GGG GGA CCG TCA GTC TTC CTC TTC CCC CCA AAA 3973  
 45 Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys  
 1125 1130 1135  
 50 CCC AAG GAC ACC CTC ATG ATC TCC CGG ACC CCT GAG GTC ACA TGC GTG 4021  
 Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val  
 1140 1145 1150

55

EP 0 896 002 A1

5 GTG GTG GAC GTG AGC CAC GAA GAC CCT GAG GTC AAG TTC AAC TGG TAC 4069  
 Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr  
 1155 1160 1165  
 10 GTG GAC GGC GTG GAG GTG CAT AAT GCC AAG ACA AAG CCG CGG GAG GAG 4117  
 Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu  
 1170 1175 1180 1185  
 15 CAG TAC AAC AGC ACG TAC CGG GTG GTC AGC GTC CTC ACC GTC CTG CAC 4165  
 Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His  
 1190 1195 1200  
 20 CAG GAC TGG CTG AAT GCC AAG GAG TAC AAG TGC AAG GTC TCC AAC AAA 4213  
 Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys  
 1205 1210 1215  
 25 GCC CTC CCA GCC CCC ATC GAG AAA ACC ATC TCC AAA GCC AAA 4255  
 Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys  
 1220 1225 1230  
 30 GGTGGGACCC GTGGGGTGG AGGGCCACAT GGACAGAGGC CGGCTCGGCC CACCCTCTGC 4315  
 CCTGAGAGTG ACCGCTGTAC CAACCTCTGT CCTACA GGG CAG CCC CGA GAA CCA 4369  
 Gly Gln Pro Arg Glu Pro  
 1235  
 35 CAG GTG TAC ACC CTG CCC CCA TCC CGG GAT GAG CTG ACC AAG AAC CAG 4417  
 Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln  
 1240 1245 1250  
 40 GTC AGC CTG ACC TGC CTG GTC AAA GGC TTC TAT CCC AGC GAC ATC GCC 4465  
 Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala  
 1255 1260 1265  
 45 GTG GAG TGG GAG AGC AAT GGG CAG CCG GAG AAC AAC TAC AAG ACC ACG 4513  
 Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr  
 1270 1275 1280 1285  
 50  
 55

5 CCT CCC GTG CTG CAT TCC GAC GGC TCC TTC TTC CTC TAC AGC AAG CTC 4561  
 Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu  
 1290 1295 1300  
 10 ACC GTG GAC AAG AGC AGG TGG CAG CAG GGG AAC GTC TTC TCA TGC TCC 4609  
 Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser  
 1305 1310 1315  
 15 GTG ATG CAT GAG GCT CTG CAC AAC CAC TAC ACG CAG AAG AGC CTC TCC 4657  
 Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser  
 1320 1325 1330  
 20 CTG TCT CCG GGT AAA TGA 4675  
 Leu Ser Pro Gly Lys  
 1335

Sequence No. 20

25 Length of sequence: 27

Type of sequence: Nucleic acid

Number of strands: Single

30 Topology: Linear

Kind of sequence: Other nucleic acid, synthetic DNA

Sequence

35 GCTCGAGCAA ACCCAGCGCA ACTACGG 27

Sequence No. 21

40 Length of sequence: 21

Type of sequence: Nucleic acid

Number of strands: Single

45 Topology: Linear

Kind of sequence: Other nucleic acid, synthetic DNA

Sequence

50 ATAGTCCCCT GATGACCATT G 21

55

Sequence No. 22

Length of sequence: 22

Type of sequence: Nucleic acid

Number of strands: Single

Topology: Linear

Kind of sequence: Other nucleic acid, synthetic DNA  
Sequence

GATGGCTTTA ATGATGTGAT TG

22

Sequence No. 23

Length of sequence: 21

Type of sequence: Nucleic acid

Number of strands: Single

Topology: Linear

Kind of sequence: Other nucleic acid, synthetic DNA  
Sequence

TGTTGGTACT TCGGCTTTCT C

21

Sequence No. 24

Length of sequence: 8

Type of sequence: Amino acid

Topology: Circular

Kind of sequence: Peptide

Sequence

Cys Ile Pro Glu Leu Ile Val Cys

1

5

Sequence No. 25

Length of sequence: 8

Type of sequence: Amino acid

Topology: Circular

Kind of sequence: Peptide

Sequence

Cys Met Arg Tyr Thr Ser Ala Cys

1 5

Sequence No. 26

Length of sequence: 8

Type of sequence: Amino acid

Topology: Circular

Kind of sequence: Peptide

Sequence

Cys Glu Trp Met Lys Arg Phe Cys

1 5

Sequence No. 27

Length of sequence: 8

Type of sequence: Amino acid

Topology: Circular

Kind of sequence: Peptide

Sequence

Cys Tyr Thr Thr Arg Leu Lys Cys

1 5

Sequence No. 28

Length of sequence: 8

Type of sequence: Amino acid

Topology: Circular

Kind of sequence: Peptide

EP 0 896 002 A1

Sequence

5 Cys Leu Arg Tyr Ser Val Pro Cys  
1 5

10 Sequence No. 29  
Length of sequence: 8  
Type of sequence: Amino acid  
15 Topology: Circular  
Kind of sequence: Peptide  
Sequence

20 Cys Ile Val Asn Arg Leu Gly Cys  
1 5

25 Sequence No. 30  
Length of sequence: 8  
Type of sequence: Amino acid  
30 Topology: Circular  
Kind of sequence: Peptide  
Sequence

35 Cys Gly Leu Gln Ala Leu Pro Cys  
1 5

40 Sequence No. 31  
Length of sequence: 8  
Type of sequence: Amino acid  
45 Topology: Circular  
Kind of sequence: Peptide  
Sequence

50 Cys Lys Leu Lys Gly Thr Met Cys  
1 5

55

## Claims

1. A chimeric protein comprising the  $\alpha$  chain or  $\beta$  chain of an integrin and the heavy chain or light chain of an immunoglobulin.
2. A chimeric protein heterodimer complex, characterized in that a chimeric protein stated in claim 1 comprising the  $\alpha$  chain of an integrin and the heavy chain or light chain of an immunoglobulin and a chimeric protein stated in claim 1 comprising the  $\beta$  chain of the integrin and the heavy chain or light chain of the immunoglobulin are associated with each other.
3. A chimeric protein heterodimer complex, according to claim 2, wherein the chimeric proteins stated in claim 1 are associated with each other in any of the following combinations (1), (2) and (3):
  - (1) An  $\alpha$  chain - immunoglobulin heavy chain -  $\beta$  chain - immunoglobulin heavy chain chimeric protein heterodimer complex, in which a chimeric protein comprising the  $\alpha$  chain of an integrin and the heavy chain of an immunoglobulin and a chimeric protein comprising the  $\beta$  chain of the integrin and the heavy chain of the immunoglobulin are associated with each other.
  - (2) An  $\alpha$  chain - immunoglobulin heavy chain -  $\beta$  chain - immunoglobulin light chain chimeric protein heterodimer complex, in which a chimeric protein comprising the  $\alpha$  chain of an integrin and the heavy chain of an immunoglobulin and a chimeric protein comprising the  $\beta$  chain of the integrin and the light chain of the immunoglobulin are associated with each other.
  - (3) An  $\alpha$  chain - immunoglobulin light chain -  $\beta$  chain - immunoglobulin heavy chain chimeric protein heterodimer complex, in which a chimeric protein comprising the  $\alpha$  chain of an integrin and the light chain of an immunoglobulin and a chimeric protein comprising the  $\beta$  chain of the integrin and the heavy chain of the immunoglobulin are associated with each other.
4. A chimeric protein or a chimeric protein heterodimer complex, according to any one of claims 1 through 3, wherein the  $\alpha$  chain of an integrin is  $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 3$ ,  $\alpha 4$ ,  $\alpha 5$ ,  $\alpha 6$ ,  $\alpha 7$ ,  $\alpha 8$ ,  $\alpha 9$ ,  $\alpha v$ ,  $\alpha L$ ,  $\alpha M$ ,  $\alpha X$ ,  $\alpha Ib$  or  $\alpha E$ .
5. A chimeric protein or a chimeric protein heterodimer complex, according to any one of claims 1 through 3, wherein the  $\beta$  chain of an integrin is  $\beta 1$ ,  $\beta 2$ ,  $\beta 3$ ,  $\beta 4$ ,  $\beta 5$ ,  $\beta 6$ ,  $\beta 7$  or  $\beta 8$ .
6. A chimeric protein heterodimer complex, according to claim 2 or 3, wherein the  $\alpha$  chain of an integrin is  $\alpha 4$  or  $\alpha 2$  and the  $\beta$  chain is  $\beta 1$ .
7. A chimeric protein or a chimeric protein heterodimer complex, according to any one of claims 1 through 3, wherein the chimeric protein comprising the  $\alpha 4$  of an integrin and the heavy chain of an immunoglobulin is identified as the amino acid sequence of sequence No. 1.
8. A chimeric protein or a chimeric protein heterodimer complex, according to any one of claims 1 through 3, wherein the chimeric protein comprising the  $\alpha 2$  of an integrin and the heavy chain of an immunoglobulin is identified as the amino acid sequence of sequence No. 19.
9. A chimeric protein or a chimeric protein heterodimer complex, according to any one of claims 1 through 3, wherein the chimeric protein comprising the  $\beta 1$  of an integrin and the heavy chain of an immunoglobulin is identified as the amino acid sequence of sequence No. 2.
10. A DNA coding for a chimeric protein stated in claim 1.
11. A DNA coding for a chimeric protein stated in claim 1, wherein the  $\alpha$  chain of an integrin is  $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 3$ ,  $\alpha 4$ ,  $\alpha 5$ ,  $\alpha 6$ ,  $\alpha 7$ ,  $\alpha 8$ ,  $\alpha 9$ ,  $\alpha v$ ,  $\alpha L$ ,  $\alpha M$ ,  $\alpha X$ ,  $\alpha Ib$  or  $\alpha E$ .
12. A DNA coding for a chimeric protein stated in claim 1, wherein the  $\beta$  chain of an integrin is  $\beta 1$ ,  $\beta 2$ ,  $\beta 3$ ,  $\beta 4$ ,  $\beta 5$ ,  $\beta 6$ ,  $\beta 7$  or  $\beta 8$ .
13. A DNA, according to claim 11, which is identified as the nucleotide sequence of sequence No. 1 or 19.
14. A DNA, according to claim 12, which is identified as the nucleotide sequence of sequence No. 2.



15. A recombinant vector, wherein a DNA stated in claim 10 is functionally linked to an expression control sequence.
16. A recombinant vector, wherein a DNA stated in claim 11 is functionally linked to an expression control sequence.
- 5 17. A recombinant vector, wherein a DNA stated in claim 12 is functionally linked to an expression control sequence.
18. A recombinant vector, wherein a DNA stated in claim 13 is functionally linked to an expression control sequence.
19. A recombinant vector, wherein the DNA stated in claim 14 is functionally linked to an expression control sequence.
- 10 20. An animal cell, comprising being transfected simultaneously by a recombinant vector in which a DNA coding for a chimeric protein comprising the  $\alpha$  chain of an integrin and the heavy chain or light chain of an immunoglobulin is functionally linked to an expression control sequence, and a recombinant vector in which a DNA coding for a chimeric protein comprising the  $\beta$  chain of the integrin and the heavy chain or light chain of the immunoglobulin is functionally linked to an expression control sequence.
- 15 21. An animal cell, according to claim 20, which is transfected simultaneously by the recombinant vectors stated in claims 16 and 17.
- 20 22. An animal cell, according to claim 20, which is transfected simultaneously by the recombinant vectors stated in claims 18 and 19.
23. A method for producing the chimeric protein heterodimer complex stated in claim 2, comprising culturing the animal cell stated in claim 20.
- 25 24. A drug, comprising a chimeric protein or chimeric protein heterodimer complex stated in any one of claims 1 through 9.
- 25 25. A drug composition, comprising a chimeric protein or chimeric protein heterodimer complex stated in any one of claims 1 through 9.
- 30 26. A platelet substitute, comprising an isolated extracellular matrix receptor as an active ingredient.
- 35 27. A platelet substitute, according to claim 26, wherein the extracellular matrix receptor is an integrin.
- 35 28. A platelet substitute, according to claim 27, wherein the  $\alpha$  chain of an integrin is  $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 3$ ,  $\alpha 4$ ,  $\alpha 5$ ,  $\alpha 6$ ,  $\alpha 7$ ,  $\alpha 8$ ,  $\alpha 9$ ,  $\alpha v$ ,  $\alpha L$ ,  $\alpha M$ ,  $\alpha X$ ,  $\alpha IIb$  or  $\alpha E$ .
- 40 29. A platelet substitute, according to claim 27, wherein the  $\beta$  chain of an integrin is  $\beta 1$ ,  $\beta 2$ ,  $\beta 3$ ,  $\beta 4$ ,  $\beta 5$ ,  $\beta 6$ ,  $\beta 7$  or  $\beta 8$ .
- 40 30. A platelet substitute, according to claim 27, wherein the integrin is integrin  $\alpha 2 \beta 1$ .
- 45 31. A platelet substitute, according to claim 26, wherein the extracellular matrix receptor is a chimeric protein heterodimer complex comprising an extracellular matrix receptor and an immunoglobulin.
- 45 32. A platelet substitute, according to claim 31, wherein the chimeric protein heterodimer complex is a chimeric protein heterodimer complex comprising an integrin and an immunoglobulin.
- 50 33. A platelet substitute, according to claim 32, wherein the chimeric protein heterodimer complex is the chimeric protein heterodimer complex stated in claim 2.
- 50 34. A platelet substitute, according to claim 33, wherein the chimeric protein heterodimer complex is the chimeric protein heterodimer complex stated in claim 6.
- 55 35. A platelet substitute, according to any one of claims 26 through 34, wherein the extracellular matrix receptor is bound to a carrier when used.
36. A platelet substitute, according to any one of claims 26 through 35, which is hemostatic.

37. A method for testing the binding between a chimeric protein heterodimer complex stated in any one of claims 2 to 9, and a ligand or cells, comprising the steps of bringing a chimeric protein heterodimer complex comprising an integrin and an immunoglobulin, and a ligand or cells into contact with each other, to prepare a mixture, and measuring the amount of the chimeric protein heterodimer complex bound to the ligand or cells or the amount of the ligand or cells bound to the chimeric protein heterodimer complex.
38. A method for searching for a substance capable of being bound to an integrin, comprising using a chimeric protein heterodimer complex stated in any one of claims 2 through 9.
39. A substance capable of being bound to an integrin, obtained by using the method stated in claim 38.
40. A method for searching for a substance which inhibits the binding between an integrin and a ligand, comprising using the method stated in claim 37.
41. A method, according to claim 40, wherein the ligand is a fibronectin fragment identified as sequence No. 3 or a collagen.
42. A protein, peptide or low molecular weight compound which inhibits the binding between an integrin and a ligand, obtained by using the method stated in claim 40 or 41.
43. A method for measuring the amount of a ligand of an integrin, comprising using a chimeric protein heterodimer complex stated in any one of claims 2 through 9.
44. A method for identifying an extracellular matrix exposed region, comprising using a chimeric protein heterodimer complex stated in any one of claims 2 through 9.

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5





Fig. 6



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP98/00370

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> Int.Cl <sup>6</sup> C07K14/705, C12N15/12, G01N33/50		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) Int.Cl <sup>6</sup> C07K14/705, C12N15/12, G01N33/50		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) BIOSIS (DERWENT), WPI (DERWENT), GenBank/EMBL (geneseq)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO, 92/13559, A1 (PROGENICS PHARMACEUTICALS, INC.), August 20, 1992 (20. 08. 92) & AU, 9214385, A	1-25, 26-34, 37-38, 40, 41
Y	WO, 95/02421, A1 (ALKERMES INC.), January 26, 1995 (26. 01. 95) (Family: none)	1-25, 26-34, 37-38, 40, 41
Y	JP, 6-87899, A (Educational Foundation Fujita Gakuen), March 29, 1994 (29. 03. 94) & EP, 466505, A2 & US, 5475100, A	1-25, 26-34, 37-38, 40, 41
Y	Nucleic Acids Res., Vol. 10, No. 13 (1982) J.W. Ellison et al., "The nucleotide sequence of a human immunoglobulin C <sub>γ</sub> gene" p.4071-4079	1-25, 26-34, 37-38, 40, 41
Y	Cell, Vol. 29, No. 2 (1982) N. Takahashi et al., "Structure of human immunoglobulin gamma genes: implications for evolution of gene family", p.671-679	1-25, 26-34, 37-38, 40, 41
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search June 8, 1998 (08. 06. 98)		Date of mailing of the international search report June 16, 1998 (16. 06. 98)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP98/00370

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EMBO J., Vol. 8, No. 5 (1989) Y. Takada et al., "The primary structure of the $\alpha^4$ subunit of VLA-4: homology to other integrins and a possible cell-cell adhesion function", p.1361-1368	1-25, 26-34, 37-38, 40, 41
Y	Mol. Immunol., Vol. 32 (1995) M.C. Szabo et al., "Identification of two variants of the human integrin $\alpha$ , subunit", p.1453-1454	1-25, 26-34, 37-38, 40, 41
Y	J. Cell Biol. Vol. 105 (1987) W.S. Argraves et al., "Amino acid sequence of the human fibronectin receptor", p.1183-1190	1-25, 26-34, 37-38, 40, 41
Y	US, 5516634, A (Newman P.J.), May 14, 1996 (14. 05. 96) (Family: none)	1-25, 26-34, 37-38, 40, 41
Y	J. Cell Biol., Vol. 109 (1989) Y. Takada et al., "The primary structure of the VLA-2/collagen receptor $\alpha^1$ subunit (platelet GPIa): homology to other integrins and the presence of a possible collagen-binding domain", p.397-407	1-25, 26-34, 37-38, 40, 41
Y	J. Clin. Invest., Vol. 92 (1993) S. Santoso et al., "The human platelet alloantigens Br <sup>a</sup> and Br <sup>b</sup> are associated with a single amino acid polymorphism on glycoprotein Ia (Integrin subunit $\alpha 2$ )", p.2427-2432	1-25, 26-34, 37-38, 40, 41
X/Y	JP, 7-500721, A (The Regents of the University of California), January 26, 1995 (26. 01. 95) & WO, 92/12236, A1	39/37-38, 43, 44
X/Y	JP, 5-505179, A (La Jolla Cancer Research Foundation), August 5, 1993 (05. 08. 93) & WO, 91/09874, A & EP, 507836, A & US, 5169930, A	39/37-38, 43, 44
Y	JP, 5-502228, A (Scripps Clinic & Research Foundation), April 22, 1993 (22. 04. 93) & WO, 91/07977, A & EP, 502124, A & US, 5196511, A	26-36, 42

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**